

# Quantum Model for the Direct Currents of Becker

M. Pitkänen

Email: [matpitka@luukku.com](mailto:matpitka@luukku.com).

[http://tgdtheory.com/public\\_html/](http://tgdtheory.com/public_html/).

April 28, 2013

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Connection between laser induced healing, acupuncture, and association of DC currents with the healing of wounds</b>	<b>3</b>
2.1	Observations relating to CNS . . . . .	3
2.2	Observations relating to regeneration . . . . .	4
2.3	Gene activation by electrostatic fields? . . . . .	5
2.4	A TGD based model for the situation . . . . .	6
<b>3</b>	<b>Quantum model for effective semiconductor property</b>	<b>10</b>
3.1	Basic model . . . . .	11
3.2	Explicit form of Schrödinger equation . . . . .	12
3.3	Semiclassical treatment . . . . .	13
3.4	Possible quantum biological applications . . . . .	13
3.5	The effects of ELF em fields on vertebrate brain . . . . .	15
3.6	Effects of 50 Hz magnetic fields on living matter . . . . .	16
3.7	The effects of polarized light on living matter . . . . .	16
3.8	Support for the proposed interaction mechanism of em radiation fields with flux tubes	17
<b>4</b>	<b>A model for remote gene expression based on Becker currents</b>	<b>18</b>
4.1	The analogy with ordinary computer . . . . .	19
4.2	Hierarchy of Planck constants and hierarchical addressing . . . . .	19
4.3	DNA supra currents and activation of genes by Becker mechanism . . . . .	19
<b>5</b>	<b>DNA, speech, music, and ordinary sound</b>	<b>20</b>
5.1	Basic picture . . . . .	21
5.2	Phonons and photons In TGD Universe . . . . .	21
5.3	What distinguishes speech and music from sounds without meaning? . . . . .	22
5.4	Are speech and music quantum duals like position and momentum? . . . . .	23

## Abstract

Motivated by a need to understand his experimental findings, Robert Becker has proposed that living matter behaves as a semiconductor in a wide range of length scales. Direct currents flowing only in a preferred direction would be essential for the functioning of living matter in this framework.

One of the key assumptions of TGD inspired theory of living matter is that various currents, even ionic currents, are quantal currents. The possibility considered mostly hitherto is that the quantal currents are Josephson currents assignable to cell membrane regarded as a Josephson junction. However, if one accepts Josephson currents, it is difficult to explain why the quantal direct currents would not be present.

The TGD inspired model for nerve pulse generation assumes that ionic currents through the cell membrane in the resting state are probably Josephson currents. If this is the case, the situation is automatically stationary and dissipation is small as various anomalies suggest. During nerve

pulse the situation could of course change and both ohmic and non-dissipating quantal direct currents could be present.

One can criticize the hypothesis about quantal DC currents since the Compton length of ions for the ordinary value of Planck constant is so small that magnetic flux tubes carrying the current through the membrane look rather long in this length scale. Therefore either Planck constant should be rather large or one should have non-ohmic quantum counterparts of a direct currents in the case of ions and perhaps also protons in the case of neuronal membrane: electronic and perhaps also protonic currents could be still Josephson currents. This would conform with the low dissipation rate.

In this article the results related to laser-induced healing, acupuncture, and DC currents are discussed. The key question is whether these direct currents are actually quantal (and thus non-dissipating) currents and whether they could be universal in living matter. A TGD inspired model for quantal direct currents is proposed and its possible implications for the model of nerve pulse are discussed.

## 1 Introduction

Robert Becker [J2] has proposed on the basis of his experimental work that living matter behaves as a semiconductor in a wide range of length scales ranging from brain scale to the scale of the entire body. Direct currents flowing only in a preferred direction would be essential for the functioning of living matter in this framework.

One of the basic ideas of TGD inspired theory of living matter is that various currents, even ionic currents, are quantal currents and do not therefore dissipate. The first possibility is that they are Josephson currents associated with Josephson junctions but already this assumption more or less also implies quantal versions of direct currents.

The TGD inspired model for nerve pulse [K6] assumes that during resting state the ionic currents through the cell membrane are Josephson currents. If this is the case, the situation is automatically stationary and dissipation is small as various anomalies suggest. During nerve pulse the situation could of course change and the both dissipating ohmic currents and non-dissipating direct currents could be present. One can criticize the hypothesis about quantal DC currents since the Compton length of ions for the ordinary value of Planck constant is so small that magnetic flux tubes carrying the current through the membrane look rather long in this length scale. Therefore either Planck constant should be rather large or one should have a non-ohmic quantum counterpart of a direct current in the case of ions and perhaps also protons in the case of neuronal membrane: electronic and perhaps also protonic currents could be still Josephson currents. This would conform with the low dissipation rate.

In the sequel the results related to laser-induced healing, acupuncture and DC currents are discussed first. The obvious question is whether these direct currents are actually electric currents (in TGD framework also so called  $Z^0$  charge could give rise to currents [K1]) and whether they could be universal in living matter. A TGD inspired model for quantal direct currents is proposed and its possible implications for the model of nerve pulse are discussed.

The model is based on sequence of questions, observations, and ideas. Why Becker's DC currents are needed? How do they relate to metabolism? How meridians and acu-points are realized in TGD Universe? What is the quantum physical correlate for "chi"? The proposed answer is that Becker's DC current consist of electron Cooper pairs identifiable as "chi", and that they are needed to compensate the leakage of electrons in electron transport cycle associated with both photosynthesis and cell respiration and providing energy for the high energy phosphate bond of ATP. Acu-points in negative potential would correspond to local stores of electron Cooper pairs compensating the electron leakage in environment. Electronic Cooper pairs are necessary for the metabolic activity: if the sign of DC potential changes, metabolism slows down and consciousness is lost. The sleep-awake rhythm and also other bio-rhythms are expected to correlate with the period change of the sign of the potential. The effects of laser light inspire the idea that also dark photons (transforming to biophotons of same energy) can take the role of solar photons and that also the emission of negative energy dark photons by electron Cooper pairs or electrons allows realization of remote metabolism by quantum credit card mechanism. Effective semiconductor property results if the transfer of electrons takes place from mitochondrial cell member space-time sheet to lipid layer space-time sheets as electron Cooper absorbs the energy liberated by oxidation of say sugar or dark photon or sends a negative energy dark photon to energy storage.

Whether the model for quantum direct currents is consistent with the proposed vacuum extremal property of the cell membrane [K6] remains an open question but both options explain the special role of  $Ca^{++}$  currents and the current of  $Na^+$  Cooper pairs in the generation of nerve pulse, as in would take place in TGD inspired model. Many-sheeted space-time allows space-time sheets which can be either almost-vacuum extremals or far from vacuum extremals - as well as space-time sheets for which the effective Planck constant is so large that both electronic and protonic Josephson currents become possible. Various pumps and channels could actually correspond to magnetic flux tubes along which various ionic supra currents or even Josephson currents can flow. The condition that both electronic and protonic supra currents are possible in the same length scale leads to the hierarchy of effective Planck constants coming approximately as powers of  $m_p/m_e \simeq 2^{11}$  proposed originally as a general truth. Radiation at Josephson frequency serves as a signature for Josephson currents.

A TGD inspired quantum model for the direct currents of Becker as direct quantum currents is developed and shown to be consistent with what is known about nerve pulse generation. The model of nerve pulse based on this model is discussed in [K6].

## **2 Connection between laser induced healing, acupuncture, and association of DC currents with the healing of wounds**

The findings of Robert Becker (the book "Electromagnetism and Life" by Becker and Marino can be found from web [J2]) meant a breakthrough in the development of bioelectromagnetics. One aspect of bioelectromagnetic phenomena was the discovery of Becker that DC currents and voltages play a pivotal role in various regeneration processes. Why this is the case is still poorly understood and Becker's book is a treasure trove for anyone ready to challenge existing dogmas. The general vision guiding Becker can be summarized by a citation from the introduction of the book.

*Growth effects include the alteration of bone growth by electromagnetic energy, the restoration of partial limb regeneration in mammals by small direct currents, the inhibition of growth of implanted tumors by currents and fields, the effect upon cephalocaudal axis development in the regenerating flatworm in a polarity-dependent fashion by applied direct currents, and the production of morphological alterations in embryonic development by manipulation of the electrochemical species present in the environment. This partial list illustrates the great variety of known bioelectromagnetic phenomena.*

*The reported biological effects involve basic functions of living material that are under remarkably precise control by mechanisms which have, to date, escaped description in terms of solution biochemistry. This suggests that bioelectromagnetic phenomena are fundamental attributes of living things, ones that must have been present in the first living things. The traditional approach to biogenesis postulates that life began in an aqueous environment, with the development of complex molecules and their subsequent sequestration from the environment by membranous structures. The solid-state approach proposes an origin in complex crystalline structures that possess such properties as semiconductivity, photoconductivity, and piezoelectricity. All of the reported effects of electromagnetic forces seem to lend support to the latter hypothesis.*

### **2.1 Observations relating to CNS**

The following more quantitative findings, many of them due to Becker, are of special interest as one tries to understand the role of DC currents in TGD framework.

1. CNS and the rest of perineural tissue (tissue surrounding neurons also including also glial cells) form a dipole-like structure with neural system in positive potential and perineural tissue in negative potential. There is also an electric field along the neuron in the direction of nerve pulse propagation (dendrites correspond to - and axon to +) (note that motor nerves and sensory nerves form a closed loop). Also microtubules within axon carry electric field and these fields are probably closely related by the many-sheeted variants of Gauss's and Faraday's laws implying that voltages along two different space-time sheets in contact at two points are the same in a static situation.
2. A longitudinal potential along front to back in the brain with the frontal lobes in negative potential with respect to occipital lobes and with a magnitude of few mV was discovered. The

strength of the electric field correlates with the level of consciousness. As the potential becomes weaker and changes sign, consciousness is lost. Libet and Gerard observed traveling waves of potentials across the cortical layers (with speeds of about 6 m/s: TGD inspired model of nerve pulse predicts this kind of waves [K6] ). Propagating potentials were also discovered in glial cells. The interpretation was in terms of electrical currents.

3. It was found that brain injury generated positive polarization so that the neurons ceased to function in an area much larger than the area of injury. Negative shifts of neuronal potentials were associated with incoming sensory stimuli and motor activity whereas sleep was associated with a positive shift. Very small voltages and currents could modulate the firing of neurons without affecting the resting potential. The "generating" potentials in sensory receptors inducing nerve pulse were found to be graded and non-propagating and the sign of the generating potential correlated with sensory input (say increase/reduction of pressure). Standard wisdom about cell membrane has difficulties in explaining these findings.
4. The natural hypothesis was that these electric fields are accompanied by DC currents. There are several experimental demonstrations for this. For instance, the deflection of assumed DC currents by an external magnetic field (Hall effect) was shown to lead to a loss of consciousness.

## 2.2 Observations relating to regeneration

The second class of experiments used artificial electrical currents to enhance regeneration of body parts. These currents are nowadays used in clinical practice to induce healing or retard tumor growth. Note that tissue regeneration is a genuine regeneration of an entire part of the organism rather than mere simple cell replication. Salamander limb generation is one of the most studied examples. Spontaneous regeneration becomes rare at higher evolutionary levels and for humans it occurs spontaneously only in the fractures of long bones.

1. An interesting series of experiments on Planaria, a species of simple flatworm with a primitive nervous system and simple head-to-tail axis of organization, was carried out. Electrical measurements indicated a simple head-tail dipole field. The animal had remarkable regenerative powers; it could be cut transversely into a number of segments, all of which would regenerate a new total organism. The original head-tail axis was preserved in each regenerate, with that portion nearest the original head end becoming the head of the new organism. The hypothesis was that the original head-tail electrical vector persisted in the cut segments and provided the morphological information for the regenerate. The prediction was that the reversal of the electrical gradient by exposing the cut surface to an external current source of proper orientation should produce some reversal of the head-tail gradient in the regenerate. While performing the experiment it was found that as the current levels were increased the first response was to form a head at each end of the regenerating segment. With still further increases in the current the expected reversal of the head-tail gradient did occur, indicating that the electrical gradient which naturally existed in these animals was capable of transmitting morphological information.
2. Tissue regeneration occurs only if some minimum amount of neural tissue is present suggesting that CNS plays a role in the process although the usual neural activity is absent. The repeated needling of the stump had positive effect on regeneration and the DC current was found to be proportional to innervation. Hence needling seems to stimulate innervation or at least inducing formation of DC currents. Something like this might occur also in the case of acupuncture.
3. Regeneration involves de-differentiation of cells to form a blastema from which the regenerated tissue is formed. Quite early it was learned that carcinogens induce de-differentiation of cells because of their steric properties and by making electron transfer possible and that denervation induces tumor formation. From these findings Becker concluded that the formation of blastema could be a relatively simple process analogous to tumor growth whereas the regeneration proper is a complex self-organization process during which the control by signals from CNS are necessary and possibly realized in terms of potential waves.
4. Regeneration is possible in salamanders but not in frogs. This motivated Becker and collaborators to compare these situations. In an amputated leg of both salamander and frog the original

negative potential of approximately -1 mV was raised first to a positive value of about +10 mV. In the frog it returned smoothly to its original value without regeneration. In the salamander it returned over a period of three days to the original base line and then went to a much higher negative value around -20 mV (resting potential is around -70 mV) followed by a return to the original value once regeneration had occurred. Thus the large negative potential is necessary for the regeneration and responsible for the formation of blastema. Furthermore, artificial electron current also induced regeneration also in the case of the frog, even in the denervated situation. Thus the flow of electrons to the stump seems to be necessary for the formation of blastema and the difference between salamander and frog is that frog is not able to provide the needed electronic current although positive potential is present.

5. It was also learned that a so called neuroepidermal junction (NEJ) formed in the healing process of salamander stump was responsible for the regeneration in the presence of denervation. The conclusion was that the DC voltage and electronic current relevant for regeneration could be assigned the interface between CNS and tissue rather than to the entire nerve and the regeneration seemed to be a local process, perhaps a feed of metabolic energy driving self-organization. Furthermore, NEJ seemed to make possible the flow of electrons from CNS to the stump.
6. The red blood cells of animals other than mammals are complete and thus possess nuclei. Becker and collaborators observed that red blood cells also dedifferentiated to form blastemas. Being normally in a quiescent state, they are ideal for studying de-differentiation. It was found that the electric current acted as a trigger at the level of cell membrane inducing de-differentiation reflected as an increased amount of mRNA serving as marker of gene expression. Also pulsed magnetic field was found to trigger the de-differentiation, perhaps via induced electric field. By the way, the role of the cell membrane fits nicely with the TGD inspired view about DNA-cell membrane system as topological quantum computer with magnetic flux tubes that are assumed to connect DNA and cell membrane and serve as braid strands in TGD inspired model of DNA as topological quantum computer [K2].
7. The experiments of Becker and collaborators support the identification of the charge carriers of DC currents responsible for the formation of the stump's large negative potential as electrons. The test was based on the different temperature dependence of electronic and protonic conductivities. Electronic conductivity increases with temperature and protonic conductivity decreases and an increase was observed.

### 2.3 Gene activation by electrostatic fields?

The basic question concerns the method of activation. The discovery of chemists Guido Ebner and Guido Schuerch [J1] raises the hope that these ideas might be more than over-active imagination and their work also provides a concrete proposal for the activation mechanism. Ebner and Schuerch studied the effect of electrostatic fields on the growth and morphogenesis of various organisms. Germ, seeds, or eggs were placed between conducting plates creating an electric field in the range .5-2 kV/m: note that the Earth's electric field is in the range .1 – 4 kV/m and of the same order of magnitude.

The outcome was rather surprising and in the year 1989 their employer Ciba Geigy (now Novartis) applied for a patent "Method of enhanced fish breeding" [J1] for what is called Ciba Geigy effect. The researchers describe how fishes (trouts) develop and grow much better, if their eggs have been conditioned in an electrostatic field. The researchers also reported [J1] that the morphology of the fishes was altered to what seems to represent an ancient evolutionary form: this was not mentioned in the patent.

The chemists founded their own Institute of Pharmaceutical Research near Basel, where Guido Ebner applied for another very detailed patent, which was never granted. In the patent he describes the effect of electrostatic fields on several life forms (cress, wheat, corn, fern, micro-organisms, bacteria) in their early stage of development. A clear change in the morphogenesis was observed. For instance, in one example fern had all sort of leaves in single plant apparently providing a series of snapshots about the evolution of the plant. The evolutionary age of the first leaf appeared to be about 300 million years whereas the last grown-up leaf looked close to its recent form.

If one takes these finding seriously, one must consider the possibility that the exposure to an electrostatic field can activate passive genes and change the gene expression so that older morphologies

are expressed. The activation of not yet existing morphologies is probably more difficult since strong consistency conditions must be satisfied (activation of program requires activation of a proper hardware). This would suggest that genome is a kind of archive also containing also older genomes even potential genomes or that topological quantum computer programs [K2] determine the morphology to a certain extent and that external conditions such as electric fields determine the self-organization patterns characterizing these programs.

It is known that the developing embryo has an electric field along the head-tail axis and that this field plays an important role in the control of growth. These fields are much weaker than the fields used in the experiment. p-Adic length scale hierarchy however predicts an entire hierarchy of electric fields and living matter is indeed known to be full of electret structures. The strength of the electric field in some p-adic length scale related to DNA might somehow serve as the selector of the evolutionary age. The recapitulation of phylogeny during ontogeny could mean a gradual shift of the activated part of both genome and "memone" (as a genetic analog of genome: for a proposal of memetic code see [K3]), perhaps assignable to topological quantum computer programs realized as braidings, and be controlled by the gradually evolving electric field strength.

The finding that led Ebner to his discovery was that it was possible to "wake up" ancient bacteria by an exposure to an electrostatic field. The interpretation would be in terms of loading of metabolic batteries. This would also suggest that in the case of primitive life forms like bacteria the electric field of the Earth has served as metabolic energy source whereas in higher life forms endogenous electric fields have taken the role of Earth's electric field.

## 2.4 A TGD based model for the situation

On the basis of these observations one can try to develop a unified view about the effects of laser light, acupuncture, and DC currents. It is perhaps appropriate to start with the following - somewhat leading - questions inspired by a strong background prejudice that the healing process - with control signals from CNS included - utilises the loading of many-sheeted metabolic batteries by supra currents as a basic mechanism.

The first series questions, observations, and ideas relates to the connection of DC currents with metabolism and ordinary biochemistry.

1. How the DC currents relate to metabolism and ordinary biochemistry? For what purpose they are needed? The crucial point is that the energy of order 1 meV gained by electron in the electric field is much below the metabolic energy quantum and also thermal energy so that the interpretation as metabolic energy quantum does not look promising. This forces to consider the possibility that the basic role of electric field is to drive electrons to where they are needed, say wounded part of tissue in positive potential and thus attracting electrons. Electrons are indeed needed by the electron transport cycle appearing in both photosynthesis and cell respiration since the transport cycle induces leakage of electrons due to the formation of ROS (reactive oxygen species) such as  $O_2^-$ . The purpose of electronic Becker currents would be therefore the re-establishment of metabolism.

The change of the sign of the Becker potential to positive induce a loss of electrons and reduced metabolism. This could explain why consciousness is lost when the sign of Becker potential is changed or electrons are deviated by Hall effect. Wound damages the connections of the tissue to the organism and the transfer of electrons compensating for leaked electrons is prevented since Becker potential changes sign. The regeneration induced by an artificial Becker potential of correct sign would induce healing by re-establishing the electron feed.

2. The role of electrons is to provide energy giving rise to the formation of ATP in cell respiration and photosynthesis. Electrons themselves receive their energy either from the oxidation of molecules or from solar photons.

The experimental support for the role of biophotons in living matter is accumulating and a natural question concerns their role in metabolism. In TGD framework dark photons with large value of  $\hbar_{eff}$  with energy of visible photon can transform to ordinary photons of same energy with some - presumably rather small - probability, and would be interpreted as biophotons. Could dark photons take the role of solar photons and provide in some situations energy to the electrons in the electron transport cycle? This would mean a non-conventional non-local

mechanism of metabolism. The effects of laser light on tissue suggest that laser light indeed takes the role of solar light and feed energy to the electron transport cycle transforming it to the energy of high energy phosphate bond of ATP.

One can consider also the possibility quantum credit card mechanism (remote metabolism) which could be at work in some the situations when chemical metabolic energy sources are absent. Damaged tissue might define this kind of exceptional situation. This brings in mind the strange ability of plants suffering under-nutrition to attract insects responsible for their pollination observed by Callahan, who has also reported that plants and insects communicate using infrared light which according to his findings serves as a sensor input in insect olfaction [I1]: also in this case quantum credit card mechanism building magnetic flux tube bridges guiding the insects to the plant might be at work. The electrons which have gained 1 meV energy during travel along pairs formed by MEs and parallel magnetic flux tubes (meridians), could send negative energy dark photons with energy of order  $\sim .5$  eV to gain same positive energy allowing to get over the semiconductor junction after they have arrived to the damaged tissue. These negative energy photons would be absorbed by a metabolic energy store (ATP in mitochondria transforming to ADP) in the healthy part of the organism.

3. The original model for the charging of the metabolic batteries and for effective semi-conductor junction assumed that the electrons of supra current are transferred to smaller space-time sheets. For ground state electrons this requires energy which is at least the difference of zero point kinetic energies of electron at the two space-time sheets. This energy should be of the order of fundamental metabolic energy quantum of about  $\sim .5$  eV.

For Cooper pairs of electrons the sheet should correspond to p-adic length scale of order  $L(k_e = 149) = 10$  nm, the thickness of lipid layer of cell membrane. For single proton corresponding scale would be  $L(k_p = 139) \simeq 2^{-12}L(151)$  from  $m_p/2m_e \simeq 2^{10}$  and  $E_{0,p}/E_{0,2e} = (2m_e/m_p) \times (L(k_e)/L(k_p))^2 = (2m_e/m_p) \times 2^{k_e - k_p} \sim 1$ .

This suggests that electron Cooper pair is kicked to a smaller space-time sheet assignable to a mitochondrial lipid layer having  $k_e = 149$ . The larger space-time sheet could be that of cell membrane with  $k = 151$ . For protons the zero point kinetic energies at these space-time sheets are by a factor  $2m_e/m_p$  lower and of the order of  $\sim .5$  meV. This happens to be of the same order of magnitude as the energy gained by proton or electron in the Becker potential. May be this is not an accident.

There is also a second intriguing quantitative co-incidence. In the absence of an action potential, acetylcholine vesicles spontaneously leak into the synaptic junction and cause very small depolarizations in the postsynaptic membrane known as miniature end plate potentials (mEPSP) of magnitude  $\sim .5$  mV (see [http://en.wikipedia.org/wiki/End-plate\\_potential](http://en.wikipedia.org/wiki/End-plate_potential)). These potentials are too small to generate action potential but together they can sum up to the needed action potential. Maybe the interpretation in terms of proton kicked to lipid layer space-time sheet might make sense.

4. The re-charging mechanism should relate directly to  $ADP \rightarrow ATP$  process occurring during electronic transport cycle in mitochondrial membrane. The connection with metabolism forces to ask how the formation of high energy phosphate bond in the addition of phosphate to ADP relates to the transfer of electrons to smaller space-time sheet. Somehow the energy of electrons must go to the formation of this bond: perhaps the dropping of electron back to larger space-time sheet transfers the energy to the high energy phosphate bond.
5. Acupuncture and the application of DC current are known to induce the generation of endorphins. Do endorphins contribute to well-being by reducing the pain? In TGD framework the deeper level interpretation of metabolism is as a provider of negentropic entanglement in turn giving rise to well-being. Are endorphins kind of negentropy packets or just conscious signals about the improved situation?

Second series of questions, observations, and ideas relates to the meridians, acu points, and "chi".

1. A permanent potential difference of same sign between head and tail could mean an accumulation of positive and negative charges to the ends of the of the system if only electron currents are

present. If both electron and proton currents with opposite directions are present, there is no accumulation of charge but there is an accumulation of protons and electrons. Probably there exists a pumping mechanism forcing the electrons (and possibly also protons) to move against the potential gradient from the tail back to the head. This however requires metabolic energy and the simplest source of this energy would be just the energy of electrons otherwise used to generate ATP. If so, the leakage would not be an unavoidable dissipative effect but a manner to avoid charge accumulation.

If the pumping mechanism is not at work, this situation cannot continue for ever and the sign of the potential difference must eventually change and induce loss of consciousness. The simplest possibility is that the potential difference changes sign rhythmically. A natural question is whether the sleep-awake rhythm is unavoidable and corresponds to the oscillatory behavior of the head-to-tail voltage.

2. "Chi" would correspond electrons or their Cooper pairs in this picture. Abnormal chi flow (reduced flow, flow in wrong direction, accumulation of chi) would cause various problems including also insomnia in which too much electron charge tends to accumulate.
3. What is the nature of acupuncture meridians, what kind of currents flow along them, and why are they not directly observed? The most natural identification in TGD Universe would be in terms of magnetic flux tubes accompanied by parallel massless extremals (MEs) making possible also the propagation of dark photons used for control purposes and perhaps even in metabolism as already discussed. Dark currents along pairs of MEs and magnetic flux tubes are ideal for the transfer of particles and energy.

If the length of the superconducting "wire" is long in the scale defined by the appropriate quantum scale proportional to  $\hbar$ , the classical picture makes sense and charge carriers can be said to accelerate and gain energy  $ZeV$ . For large values of  $\hbar_{eff}$  an oscillating Josephson current would be in question. Since Becker currents are associated with CNS, it would be natural to associate the meridians with neural pathways although this assignment is not necessary. Magnetic flux tube system defined kind of magnetic circulation which could serve as a template for the neural pathways. The transfer of energy with minimal dissipation would explain why a semiconductor like property is needed and why acupuncture points have a high conductivity value.

4. What about acu-points? Acu points are known to be in negative potential normally. This suggests that the density of electrons or their Cooper pairs at them is higher than elsewhere in the meridian. Could they serve as kind of electron stores providing electrons to their environment to compensate for losses caused by ROS. This would make possible higher metabolic activity in presence of nutrient molecules since the rate for the electron transform cycle should be proportional to the density of energizable electrons, "chi".

When the potential of the acu-point is reduced or become positive, under-nutrition follows. This should relate to various symptoms like pain at acupuncture points. Acupuncture needle as an electronic conductor would develop a charge distribution with a concentration of electrons to the acu-point, and would re-establish the metabolic activity. Pain would be signature of lack of negentropic entanglement and positive/negative coloring of emotions and sensations would quite generally correlate with the amount of negentropic entanglement.

5. Nanna Goldman et al have provided empirical evidence [I3] for the expectation that the healing effect of the acupuncture involves metabolism (see the popular article in Sciencedaily [I2]).

The group has found that adenosine is essential for the pain killing effects of acupuncture. For mice with a normal adenosine level acupuncture reduced dis-comfort by two-thirds. In special "adenosine receptor knock-out mice" acupuncture had no effect. When adenosine was turned on in the tissues, the discomfort was reduced even in the absence of acupuncture. During and after an acupuncture treatment, the level of adenosin in tissues near the needles was 24 times higher than before the treatment. In the abstract of the article it is stated that it is known for long time that acupuncture generates signals which induce brain to generate natural pain killing endorphins but that also adenosine acts as a natural pain killer.



Adenosine is the basic building block of AXP,  $X=M,D,T$  (adenosin-X-phosphate,  $X=mono,di,tri$ ). Therefore the findings suggest that the flow of electrons from the needle to acu point loads metabolic batteries by providing electrons to electron transport cycle needed to generate ATP. Adenosine could be partially generated as decay products of AXPs. Tissue itself could increase adenosine concentration to make possible its transformation to AXP utilizing electric field energy. From the popular article one cannot conclude whether the authors propose a connection with metabolism. The results are consistent with the assumption that the AXPs generated from adenosin accompany negentropic entanglement. This can occur in the scale of entire body and meridians could also make possible direct signalling with brain.

How can understand the semiconducting character of Becker's DC currents?

1. Becker assigns to the system involved with DC currents an effective semiconductor property. Could the effective semiconductor property be due the fact that the transfer of charge carriers to a smaller space-time sheet by first accelerating them in electric field is analogous to the transfer of electrons between conduction bands in semiconductor junction? If so, semiconductor property would be a direct signature of the realization of the metabolic energy quanta as zero point kinetic energies. For metabolic energy quantum of order .5 eV this however makes sense only if the electrons transferred to the smaller space-time sheet have energy slightly below the minimum energy for the transfer to the smaller space-time sheet in absence of the Becker potential. The situation would be critical and 1 mV voltage could serve as a kind of control knob.
2. Supra currents flowing along magnetic flux tubes would make possible dissipation-free loading of metabolic energy batteries. This even when oscillating Josephson currents are in question since the transformation to ohmic currents in semiconductor junction makes possible energy transfer only during second half of oscillation period. Could this be a universal mechanism applying to various stages of the regeneration process? In quantal situation the metabolic energy quanta have very precise values as indeed required.
3. The findings of Becker provide support for electronic DC currents. The Cooper pairs of electrons are indeed the best candidates for the carriers of supra current by their small mass. In the minimal situation the currents defined by leaked electrons moving against potential gradient (utilising the energy used otherwise to generate ATP) could compensate the Becker currents and give rise to closed current loops without charge accumulation. If the electronic DC currents observed by Becker are much stronger than needed to compensate for the local electron leakage, a larger metabolically driven return current is needed to guarantee local charge neutrality. These currents seem to be assignable to CNS: maybe the two electron currents could be associated with sensory and motor pathways. An interesting question whether sympathetic-parasympathetic dichotomy also relates to electron currents in opposite directions.
4. Could also dark protonic and even ionic DC currents be present and running along their own flux tubes and perhaps defining cyclotron Bose-Einstein condensates? How large the scale of flux tubes can be: could it be much larger than that of biological body (by simple argument magnetic body should have layers with even size scale of Earth). What is the possible connection with cell respiration? When single ATP is generated, three protons are pumped through the mitochondrial membrane utilising the energy liberated in electron transport cycle. This does not however require protonic currents in longer scales.
5. In regeneration process NEJs are formed. They could consist of pairs of MEs and magnetic flux tube mediating the electronic DC current during blastema generation and regeneration proper during which also control signals from CNS would be present. Since NEJs seems to resemble cell membranes in some respects, the ideas inspired by the model of cell membrane and DNA as tqc can be used. The model for nerve pulse and the model for DNA as topological quantum computer suggest that dark ionic currents flowing along magnetic flux tubes characterised by a large value of the effective Planck constant are involved with both meridians and NEJs. Magnetic flux tubes can act as DC current wires or Josephson junctions generating oscillatory supra currents of ions and electrons. Also for large values of the effective Planck constant meridians look short in the relevant dark length scale and could act as Josephson junctions carrying oscillatory Josephson currents.

One can raise also questions about the relationship between DC currents and de-differentiation.

1. Could cell de-differentiation be caused by the presence of Becker's DC current? Also acupuncture is known to induce de-differentiation. Could the mere ability to charge metabolic energy batteries provided by electron feed induce de-differentiation, which manifests as an increased genetic expression? Can one see differentiation as an eliminative process forced by the reduction of the electron feed and inducing a selective reduction of gene expression? If this were the case, de-differentiation could be induced by a feed of surplus electrons to the system using either external electron current or additional electric field. Local electron density would correlate negatively with the degree of differentiation.
2. In this framework it might be possible to understand the claimed effects of external electric fields on the development of plants and fishes. In this case rejuvenation means return to the earlier evolutionary stages. Maybe ontogeny-recapitulates-phylogeny principle might allow to understand this if genome in some sense contains archive about earlier stages of evolution. This archive might be virtual and realised by an epigenetic mechanism selecting different patterns of gene expression using the same genome.

If this is the case, the density of electrons or their Cooper pairs - "chi" - possessed by the cell would serve as a measure for the biological age of the cell and the meridian system feeding "chi" would serve as a rejuvenating agent with respect to gene expression. The average density of dark electrons would serve as a measure for the age of cell: the larger the density the higher the metabolic activity and the lower the biological age.

### 3 Quantum model for effective semiconductor property

Becker [J2] summarizes his findings by stating that living matter is an effective semiconductor. There are pairs of structures in positive and negative potential in various scales and the current between the plates of this effective capacitor flows when above some minimum potential difference. The current flows from positive to negative pole and could be an electron current. Also proton current in the opposite direction can be considered but the electron current is experimentally favored. For instance consciousness is lost when a magnetic field is used to deflect the current.

In TGD framework natural carriers of these currents would be magnetic flux tubes also carrying electric fields. A very simple deformation of the imbedding of a constant longitudinal magnetic field also gives longitudinal electric field. With a slight generalization one obtains helical electric and magnetic fields. A crucial difference is that these currents would be quantal rather than ohmic currents even in the length scale of the biological body and even longer scales assignable to the magnetic body.

The following argument allows us to understand the physical situation.

1. A precise everyday analogy is vertical motion in the gravitational field of the Earth between surface and some target at given height  $h$ . If the kinetic energy is high enough, the particle reaches the target. If not, the particle falls back. In the quantum case one expects that the latter situation corresponds to a very small probability amplitude at the target (tunneling to classically forbidden kinematic region).
2. Now the electric field replaces the gravitational field. Suppose that the classical electric force experienced by the particle is towards the capacitor plate taking the role of the surface of Earth. Below critical field strength the charged particle cannot reach the target classically and quantum mechanically this occurs only by tunneling with vanishingly small probability.
3. Particles with opposite value of charge experience a force which accelerates them and classically they certainly reach the second plate. What happens in a quantum situation? It seems that this situation is essentially identical with the first one: one has linear potential in finite interval and wave functions are localized in this range. One can equivalently regard these states as localized near the second capacitor plate.
4. A good analogy is provided by atoms: classically the electron would end down at the nucleus but quantization prevents this. One can imagine also now stationary solutions for which the electric currents for individual charges vanish at the plates although classically there would be

a current in another direction. Also quantum mechanically non-vanishing conserved current is possible: all depends on boundary conditions.

### 3.1 Basic model

Consider now the situation at more quantitative level.

1. One can assign complex order parameters  $\Psi_k$  to various Bose-Einstein condensates of supra phases and obey Schrödinger equation

$$i\partial_t\Psi_k = \left(-\frac{\hbar^2}{2m_k}\partial_z^2 + q_k Ez\right)\Psi_k . \quad (3.1)$$

Here it is assumed that the situation is effectively one-dimensional.  $E$  is the value of constant electric field.

2. The Schrödinger equation becomes non-linear, when one expresses the electric field in terms of the total surface charge density associated with the plates of effective capacitor. In absence of external electric field it is natural to assume that the net surface charge densities  $\sigma$  at the plates are of opposite sign so that the electric field inside the capacitor is proportional to

$$\sigma = E = \sum \sigma_i = \sum_i q_i \bar{\Psi}_i \Psi_i . \quad (3.2)$$

This gives rise to a non-linear term completely analogous to that in non-linear Schrödinger equation. A more general situation corresponds to a situation in which the region interval  $[a, b]$  bounded by capacitor plates  $a$  and  $b$  belongs to a flux longer tube like structure  $[A, B]$ :  $[a, b] \subset [A, B]$ . In this case one has

$$E_{tot} = E + E_0 . \quad (3.3)$$

This option is needed to explain the observations of Becker that the local strengthening of electric field increases the electron current: this would be the case in the model to be discussed if this field has a direct opposite to the background field  $E_0$ . One could also interpret  $E$  as quantized part of the electric field and  $E_0$  as classical contribution.

3. The electric currents are given by

$$j_k = \frac{i\hbar q_k}{2m_k} \bar{\Psi}_k \partial_z^{\leftrightarrow} \Psi_k . \quad (3.4)$$

In stationary situation the net current must vanish:

$$\sum_k j_k = 0 . \quad (3.5)$$

A stronger condition is that individual currents vanish at the plates:

$$j_k = 0 . \quad (3.6)$$

It must be emphasized that this condition does not make sense classically.

### 3.2 Explicit form of Schrödinger equation

Consider now the explicit form of Schrödinger equation in a given electric field.

1. The equation is easy to solve by writing the solution ansatz in polar form (the index  $k$  labelling the charge particle species will be dropped for notational convenience).

$$\Psi = R(a\exp(iU) + b\exp(-iU))\exp(-iE_n t) \quad (3.7)$$

For real solutions current vanishes identically and this is something which is not possible classically. It is convenient to restrict the consideration to stationary solutions, which are energy eigen states with energy value  $E_n$  and express the general solution in terms of these.

2. The Schrödinger equation reduces with the change of variable

$$\begin{aligned} z &\rightarrow \frac{(z - z_0)}{z_1} \equiv x , \\ z_0 &= \frac{E_n}{qE} , \quad z_1 = \left(\frac{\hbar^2}{2mqE}\right)^{1/3} . \end{aligned} \quad (3.8)$$

to

$$(\partial_x^2 + x)\Psi = 0 . \quad (3.9)$$

The range  $[0, z_0]$  for  $z$  is mapped to the range  $[-z_0/z_1, 0]$ .  $z_0/z_1$  has positive sign as is easy to verify. The value range of  $x$  is therefore negative irrespective of the sign of  $qE$ . This is the differential equation for Airy functions [B1]. Airy functions are encountered in WKB approximation obtained by linearizing the potential function:  $V(x) = ax + b + O(x^2)$ .

The change of variable leads automatically to solutions restricted near the plate where the situation is completely analogous to that in the gravitational field of the Earth. For stationary solutions a test charge in a given background field would be localized near the capacitor plate with opposite sign of charge. A strong background field could be created by charges which do not correspond to the ionic charges defining ionic currents. Electrons and protons could define this field possibly associated with flux tubes considerably longer than the distance between capacitor plates.

3. Using the polar representation  $\Psi = R\exp(iU)$  Schrödinger equation reduces to two equations

$$\begin{aligned} [(\partial_x^2 - U_x^2 + x)R] \cos(U) + [U_{xx} + 2\partial_x R \partial_x U] \sin(U) &= 0 , \\ [(\partial_x^2 - U_x^2 + x)R] \sin(U) - [U_{xx} - 2\partial_x R \partial_x U] \cos(U) &= 0 . \end{aligned} \quad (3.10)$$

Note that both  $(R, U)$  and  $(R, -U)$  represent solutions for given value of energy so that the solution can be chosen to be proportional to  $\cos(U)$  or  $\sin(U)$ . The electric current  $j$  is conserved and equal to the current at  $x = 0$  and given by

$$j = \frac{\hbar}{2m} \frac{U_x}{z_1} R^2 , \quad z_1 = \left(\frac{\hbar}{2mqE}\right)^{1/3} . \quad (3.11)$$

The current vanishes if either  $U_x$  is zero or if the solution is of form  $\Psi = R\sin(U)$ .

### 3.3 Semiclassical treatment

In semiclassical approximation the potential is regarded as varying so slowly that it can be regarded as a constant. In this situation one can write the solution of form  $Rexp(iU)$  as

$$\Psi = R_0 exp\left(\frac{i}{\hbar} \int_0^z \sqrt{2m} \sqrt{E - qEz} dz\right) = R_0 exp\left(i \int_0^x x^{1/2} dx\right) . \quad (3.12)$$

The plate at which the initial values are given can be chosen so that the electric force is analogous to gravitation at the surface of Earth. This requires only to replace the coordinate  $z$  with a new one, vanishing at the plate in question - and gives to the energies a positive shift  $E_0 = qE_0 h$ .

1. The semiclassical treatment of the equation leads to Bohr rules

$$\frac{\oint p_z dz}{\hbar} = \frac{2}{\hbar} \int_0^h p_z dz = n . \quad (3.13)$$

This gives

$$\frac{\oint p_z dz}{\hbar} = \frac{2\sqrt{2m}}{\hbar} \int_0^h \sqrt{E_n - qEz} dz = 2 \int_0^{x_0} x^{1/2} = \frac{4}{3} x_0^{3/2} = n . \quad (3.14)$$

Note that the turning point for classical orbit corresponds to  $z_{max} = E_n/qE$ .

2. One obtains

$$E_n = \frac{1}{2} \left( \frac{nqE\hbar^2}{r\sqrt{m}} \right)^{2/3} , \quad r = \int_0^1 (1-u)^{1/2} du = \frac{2}{3} . \quad (3.15)$$

The value of  $z_{max}$  is

$$z_{max} = \frac{E_n}{qE} = \frac{n^{2/3}}{2r^{2/3}} \left( \frac{\hbar^2}{qEm} \right)^{1/3} . \quad (3.16)$$

3. The approximation  $R = R_0 = constant$  can make sense only if the position of the second plate is below  $z_{max}$ . This is possible if the value of  $n$  is large enough ( $n^{2/3}$  proportionality), if the mass  $m$  of the charged particle is small enough ( $m^{-1/3}$  proportionality) raising the electron and also the proton to a special position, or if the strength of the electric field is small enough ( $E^{-1/3}$  proportionality). The value  $z_{max}$  is proportional to  $\hbar^{2/3}$  so that a phase transition increasing Planck constant can induce current flow.

### 3.4 Possible quantum biological applications

The proposed model for quantum currents could provide quantum explanation for the effective semiconductor property of Becker's DC currents.

1. The original situation would be stationary with no currents flowing. The application of an external electric field in the correct direction would reduce the voltage below the critical value and currents would start to flow. This is consistent with Becker's findings if there is a background electric field  $E_0$  with direction opposite to that of the applied field has a direction opposite to  $E_0$  so that the field strength experienced by charged particles is reduced and it is easier for them to reach the second plate.

2. Becker's DC currents appear in several scales. They are assigned with the pairs formed by CNS and perineural tissue (this includes also glia cells) and by frontal and occipital lobes. Acupuncture could involve the generation of a DC supra current. The mechanism would be essential in the healing. Also the mechanism generating qualia could involve generation of supra currents and dielectric breakdown for them. The role of the magnetic flux tubes in TGD inspired biology suggests that the mechanism could be universal. If this were the case one might even speak about a Golden Road to the understanding of living matter at the basic level.

Even the generation of nerve pulse might be understood in terms of this mechanism. One can argue that neurons have a higher evolutionary level than the system pairs to which only electron currents or electron and proton currents can be assigned. This is because the value of the effective Planck constant is higher for the magnetic flux tubes carrying the quantal ionic currents.

1. For Bose-Einstein condensate the simplest choice is  $n = 1$  at both plates. The energy eigenvalues would naturally differ by the shift  $E_0 = qE_0h$  at the two plates for a given particle type. Under these assumptions the current can flow appreciably only if the voltage is below the minimum value. This is certainly a surprising conclusion but brings in mind what happens in the case of neuronal membrane. Indeed, hyper-polarization has a stabilizing - something difficult to understand classically but natural quantum mechanically.
2. The reduction of membrane potential slightly below the resting potential generates nerve pulse. Also a phase transition increasing the value of the effective Planck constant might give rise to quantal direct currents and generate flow of ionic currents giving rise to nerve pulse. Stationary solutions are located near either capacitor plate. What comes to mind is that the nerve pulse involves a temporary change of the capacitor plate with this property.
3. If the electron and proton currents flow as direct currents, one encounters a problem. Nerve pulse should begin with direct electronic currents and be followed by direct protonic currents and only later ions should enter the game if at all. The existing model for nerve pulse however assumes that at least electrons flow as oscillating Josephson currents rather than direct quantal currents. This is quite possible and makes sense if the cell membrane thickness is small - that is comparable to electron Compton length as assumed in large  $\hbar$  model for the nerve pulse. This assumption might be necessary also for proton and would make sense if the Planck constant for protonic flux tubes is large enough. For ions the Compton length would be much smaller than the thickness of cell membrane and direct currents would be natural.

If the value of the effective Planck constant is the same for biologically important ions, direct quantum currents would be generated in definite order since in  $\hbar < z_{max}$  one has  $z_{max} \propto m^{-1/3} \propto A^{-1/3}$ . The lightest ions would start to flow first.

- (a) Nerve pulses can be generated by voltage gated channels for potassium and calcium. Voltage gated channels would correspond to magnetic flux tubes carrying electric field. For voltage gated channels  $\text{Na}^+$  ions with atomic weight  $A = 23$  and nuclear charge  $Z = 11$  start to flow first, then  $\text{K}^+$  ions with atomic weight  $A = 39$  and  $Z = 19$  follow. This conforms with the prediction that the lightest ions flow first. The nerve pulse duration is of the order of 1 millisecond at the most.
- (b) Nerve pulses can be also generated by voltage gated  $\text{Ca}^{++}$  channels. In this case the duration can be 100 ms and even longer.  $\text{Ca}$  has  $A = 40$  and  $Z = 20$ . The proper parameter is  $x = r^2/qA$ ,  $r = \hbar/h_0$ . One has

$$\frac{x(\text{Ca}^{++})}{x(\text{Na}^+)} = \left(\frac{r(\text{Ca}^{++})}{r(\text{Na}^+)}\right)^2 \times \frac{23}{2 \times 40} . \quad (3.17)$$

$r^2(\text{Ca}_{++}) \sim 2r^2(\text{Na}_+)$  would allow to compensate for the increased weight and charge of  $\text{Ca}_{++}$  ions.

4. The objection is that  $\text{Na}^+$  and  $\text{K}^+$  are not bosons and therefore cannot form Bose-Einstein condensates. The first possibility is that one has Cooper pairs of these ions. This would imply

$$\frac{x(Ca^{++})}{x(2Na^+)} = \left(\frac{r(Ca^{++})}{r(Na^+)}\right)^2 \times \frac{23}{20} .$$

$Ca^{++}$  and  $Na^+$  pair would be in very similar position for a given value of Planck constant. This is a highly satisfactory prediction. Another manner to circumvent the problem is more science fictive and assumes that the  $Na^+$  ions are exotic nuclei behaving chemically as  $Na^+$  but having one charged color bond between nucleons [K5].

It remains to be seen whether this model is consistent with the model of cell membrane as almost vacuum extremal or whether the vacuum extremal based model could be modified by treating ionic currents as direct currents. In the vacuum extremal model classical  $Z^0$  gauge potential is present and would give a contribution to the counterpart of Schrödinger equation. The ratio  $x(Ca^{++})/x(2Na^+)$  for the parameter  $x = r^2/q(A-Z)A$  (em charge  $q$  is replaced with neutron number in good approximation) equals to 1.38 and is not therefore very far from unity.

The many-sheetedness of space-time is expected to play a key role and one should precisely specify which sheets are almost vacuum extremals and which sheets are far from vacuum extremals. One expects that magnetic flux tubes are far from vacuum extremals and if voltage gated ionic channels are magnetic flux tubes, the proposed model might be consistent with the model of cell membrane as almost vacuum extremal.

### 3.5 The effects of ELF em fields on vertebrate brain

The effects of ELF em fields on vertebrate brain occur both in frequency and amplitude windows. Frequency windows can be understood if the effect occur at cyclotron frequencies and correspond to absorption of large  $\hbar$  photons. A finite variation width for the strength of magnetic field gives rise to a frequency window. The observed quantal character of these effects occurring at harmonics of fundamental frequencies leads to the idea about cyclotron Bose-Einstein condensates as macroscopic quantum phases. The above considerations support the assumption that fermionic ions form Cooper pairs.

I have tried to understand also the amplitude windows but with no convincing results. The above model for the quantum currents however suggests a new approach to the problem. Since ELF em fields are in question they can be practically constant in the time scale of the dynamics involved. Suppose that the massless extremal representing ELF em field is orthogonal to the flux tube so that the ions flowing along flux tube experience an electric force parallel to flux tube. What would happen that the ions at the flux tube would topologically condensed at both the flux tube and massless extremal simultaneously and experience the sum of two forces.

This situation is very much analogous to that defined by magnetic flux tube with longitudinal electric field and also now quantum currents could set on. Suppose that semiconductor property means that ions must gain large enough energy in the electric field so that they can leak to a smaller space-time sheet and gain one metabolic quantum characterized by the p-adic length scale in question. If the electric field is above the critical value, the quantum current does not however reach the second capacitor plate as already found: classically this is of course very weird. If the electric field is too weak, the energy gain is too small to allow the transfer of ions to smaller space-time sheet and no effect takes place. Hence one would have an amplitude window.

The amplitude window occur in widely separate ranges 1-10 V/m and around  $10^{-7}$  V/m. Of course, also other frequency ranges might be possible. Fractality and the notion of magnetic suggests a possible explanation for the widely different frequency ranges. Both p-adic length scale hypothesis and the hierarchy of Planck constants suggest that some basic structures associated with the cell membrane have fractal counterparts in a wide length scale range and correspond to binary structures. Magnetic flux tubes carrying quantal DC currents of Becker would be the most natural candidate in this respect since these currents appear in several length scales inside organism. Also the counterparts of lipid layers of cell membrane could be involved. If so, one must include to the hierarchy of amplitude windows also fields in the range corresponding to the cell membrane resting potential of about  $6 \times 10^6$  V/m. This is of course only a rough order of magnitude estimate since perturbations of these field are in order.

By fractality the most natural guess is that the voltage along the flux tube is invariant under the scale of Planck constant. This would mean that the electric field would behave as  $1/L^2 \propto 1/\hbar^2$  as a

function of the length scale characterizing the scale variant of the structure. If so the range  $E = 1 - 10$  V/m assignable also to EEG would correspond to a length scale of  $7.7 - 24 \mu\text{m}$  corresponding to cell length scale. Perhaps the direct currents run between cells layers.  $E = 10^{-7}$  V/m would in turn correspond to 7.8 cm which corresponds to size scale of human brain hemisphere (experiments were carried out for vertebrates). Could the direct quantum currents in question run between brain hemispheres along corpus callosum?

### 3.6 Effects of 50 Hz magnetic fields on living matter

The vision about the role of cyclotron Bose-Einstein condensates was inspired by the effects of ELF em fields on vertebrate brain. The magnetic field strength explaining the effects was about .2 Tesla, 2/5 of the nominal value for the strength of Earth's magnetic field.

There are also other experiments have demonstrated that oscillating electromagnetic fields have effects on living matter. In particle oscillatory magnetic fields with frequency of 50 Hz and with field strengths typically in the range .1-1 mT are used: these effects are summarized in [J4]. Even fields of order .14 Tesla are used.

It is interesting to look at the values of basic parameters associated with these fields.

1. For 50 Hz oscillation frequency the wave length  $\lambda$  is 6000 km to be compared with the radius of Earth which is 6371 km. If one takes seriously the notion of magnetic body this need not be an accident. I do not know how essential it is to have just 50 Hz frequency. The magnetic field is nearby oscillating dipole field up to distances of order  $\lambda$  and radiation field at much longer distances. Therefore the field in question is in good approximation nearby field as far as biological body is considered. For magnetic body the radiation field could dominate
2. For the endogenous magnetic field  $B_{end} = .2$  Gauss cyclotron frequencies of ions are in EEG range:  $Ca^{++}$  cyclotron frequency is 15 Hz. The scaling up to  $r = .1-1$  mT means scaling of cyclotron frequencies by a factor 5 - 50. For  $Ca^{++}$  this would give frequency range 75-750 Hz. For  $K^+$  and  $Cl^+$  ions the frequency range would be about 35-375 Hz.
3. The magnetic length  $r = \sqrt{2/eB}$  characterizing flux tube thickness for flux quantization with minimum value of flux is for  $B = .05$  mTesla equal to  $5 \mu\text{m}$ . For the fields in the range .1-1 mTesla it is in the range  $3.5 \mu\text{m} - 1.1 \mu\text{m}$ .  $2.5 \mu\text{m}$  corresponds to p-adic length scales  $L(k)$  associated with Gaussian Mersenne  $M_{G,k} = (1+i)^k - 1$ ,  $k = 167$ , and Gaussian Mersenne corresponding to  $k = 163$  would correspond to p-adic length scale  $.36 \mu\text{m}$ . .14 Tesla corresponds to magnetic length of 9.4 nm rather near to cell membrane thickness of 10 nm which corresponds to p-adic length scale  $L(151)$  assignable to Gaussian Mersenne  $M_{G,151}$ .

### 3.7 The effects of polarized light on living matter

Polarized light is known to have effects on living matter [J4]. For instance, Peter Gariaev has found that the polarized light generated by living matter sample irradiated by polarized laser light has effects on distant organism and there are even indications that genetic code might be realized in terms of radiation patterns [K8]. The quantum model for Becker currents suggest that these effects result as a modification of the voltage between the ends of magnetic flux tubes. If the flux tubes are near criticality for the generation of quantal DC currents, polarized light could be utilized both communication and control purposes whereas the acceleration in the electric fields along flux tubes would serve as a provider of metabolic energy allowing to load metabolic batteries. This process could be initiated by an electromagnetic signal inducing generation of quantal currents. The same basic mechanism could be at work also in DNA transcription, replication and other similar processes.

If the polarized low frequency radiation corresponds to a massless extremal (ME) orthogonal to the flux tube such that the polarization of the radiation is parallel to the flux tube, the voltage is affected by a contribution given by  $\Delta V = Ed$ ,  $d$  the thickness of ME. If the flux tube is near criticality to a generation of quantal currents this change of voltage could serve as a signal inducing the generation of quantal currents.

The maximal effect is obtained for the flux tubes having direction parallel to the electric polarization so that the effect is highly selective. In the case of DNA double strand the direction of flux tube changes so that the effect would be maximal on DNAs which correspond to the same angular position



on the super-coil of radius of order 10 nm formed by DNA double helix. This allows to imagine signals for which temporal variation of polarization direction means scanning of DNA.

It is known that the energy of radiation can be transformed to metabolic energy. For instance, IR light for which photons have energies of order metabolic quantum has biological effects [I5]. The mechanism could be following. Suppose that the electric field of IR photon is parallel to the flux tube which carries an electric field and is near criticality for the generation of quantal DC currents. If the direction of polarization is correct, the additional contribution to electric field induces direct current and acceleration of electrons and protons and their transfer to smaller space-time sheets and therefore loading of metabolic batteries. This could also make generation of ATP possible.

Suppose that one takes seriously the model for remote replication of DNA [K8] involving flux tubes connecting identical DNA nucleotides and that the radiation propagating along them induces quantal currents along the receiving DNA inducing replication and perhaps even transcription. The direction of polarization for the emitted radiation should be parallel to the DNA strand locally and during its travel to the target the polarization should remain orthogonal to the flux tube so that one would have what might be called polarization window. Parallel translation of the polarization vector in the induced metric suggests itself.

### 3.8 Support for the proposed interaction mechanism of em radiation fields with flux tubes

The basic prediction of the interaction mechanism is that the effects of em field with a given frequency occur only at the second half period when the direction of electric field is "correct". This prediction might be testable. In fact, there is evidence for this interaction mechanism in the case of theta waves of EEG. The memory storage occurs only at the second half of the theta wave This is discussed from different point of view in [K2].

The place coding by phase shifts was discovered by O'Reefe and Recce [J3]. In [J5, J6]. Y. Yamaguchi describes the vision in which memory formation by so called theta phase coding is essential for the emergence of intelligence. It is known that hippocampal pyramidal cells have "place property" being activated at specific "place field" position defined by an environment consisting of recognizable objects serving as landmarks. The temporal change of the percept is accompanied by a sequence of place unit activities. The theta cells exhibit change in firing phase distributions relative to the theta rhythm and the relative phase with respect to theta phase gradually increases as the rat traverses the place field. In a cell population the temporal sequence is transformed into a phase shift sequence of firing spikes of individual cells within each theta cycle.

Thus a temporal sequence of percepts is transformed into a phase shift sequence of individual spikes of neurons within each theta cycle along linear array of neurons effectively representing time axis. Essentially a time compressed representation of the original events is created bringing in mind temporal hologram. Each event (object or activity in perceptive field) is represented by a firing of one particular neuron at time  $\tau_n$  measured from the beginning of the theta cycle.  $\tau_n$  is obtained by scaling down the real time value  $t_n$  of the event. Note that there is some upper bound for the total duration of memory if scaling factor is constant.

One can say that neurons in ensemble provide a representation for the external world and the location of the rodent in the external world is represented as a firing of a neuron in this landscape. Besides this also temporal scaling down by a factor about ten is carried out so that actual event is represented as much shorter copies of it. Obviously this represents temporal fractality.

This scaling down - story telling - seems to be a fundamental aspect of memory. Our memories can even abstract the entire life history to a handful of important events represented as a story lasting only few seconds. This scaling down is thought to be important not only for the representation of the contextual information but also for the memory storage in the hippocampus. Hierarchy of Planck constants and phase transitions changing Planck constant make this story building possible.

The finding of Yamaguchi and collaborators relevant in the recent context is that the gradual phase shift occurs at half theta cycle whereas firings at the other half cycle show no correlation [J5]. The proposed model for the interaction of theta waves with flux tubes could explain this naturally. The relevant neural sub.system would be critical to the generation of quantal DC current only when the direction electric field of synchronizing theta wave generated by magnetic body is correct. Hence synchronous neural activity would be induced only at second half cycle of theta wave and firing would be random during the other half cycle.

## 4 A model for remote gene expression based on Becker currents

If one accepts the notion of magnetic body as intentional agent, the basic challenge is to understand how magnetic body realizes its intents as remote mental interactions on biological body. This model must of course apply also to the more conventional remote mental interactions such as remote realization of intent.

The hypothesis is that electromagnetic and possibly also other massless classical fields assignable to so called massless extremals are in a key role. Also cyclotron frequencies characterizing magnetic bodies play a key role. The vision is that magnetic flux sheets traverse many-sheeted DNA in various scales giving rise to a hierarchy of genomes and coherent gene expression in scales of cell, organelles, organism, and even population, and species. Hierarchy of Planck constants is in an essential role in realizing this hierarchy in terms of photons with energies above the thermal energy at physiological temperature and having spectrum of wavelengths coming as multiples  $\lambda = n\lambda_0$ ,  $n = \hbar/\hbar_0$ .

The findings of Benveniste and followers relating to water memory and homeopathy, the recent work of group led by HIV Nobelist Luc Montagnier coupling the findings with genetics and suggesting a new nanoscale realization of genetic code (see this [L1]), the work of the group of Popp with biophotons identified as decay products of large  $\hbar$  photons with visible energies (in particular dark EEG photons), and the work of Peter Gariaev and collaborators supporting remote gene expression and replication discussed here suggest that electromagnetic radiation is indeed involved [K8]. In the case of water memory and homeopathy the spectrum of cyclotron frequencies for the chemical invader characterizes it and induces immune response trying to eliminate it. I have also proposed a model for how genes coding for proteins eliminating the invader could be generated almost automatically: the model is based on the predicted realization of vertebrate genetic code in terms of dark proton states (see this [K4]). DNA would like an animal which sniffs the invaders magnetic body and automatically reacts to the smell.

The discussions with Lian Sidorov and people who have realized that new era is beginning in biology have served as a driving force in the attempts to formulate in more detail TGD inspired view about how remote mental interactions - which are basic element of the model in TGD framework - might be realized. As a matter fact, I have added to my homepage a new book summarizing briefly the recent view about quantum TGD and its applications to quantum consciousness, quantum biology, to quantum neuroscience, and to remote mental interactions with some proposals for possible tests [K7] (see this).

To start with, suppose that in the case of biological target realization of intent in the simplest situation reduces to expression of genes. This is of course a strong limitation to the type of remote mental interactions. The challenge is to develop a model for remote realization of genetic activities like replication, and transcription. For some time ago I proposed a model with Peter Gariaev [K8] but it was still too clumsy since it required too much of information transfer between the genomes of sender and receiver. Much simpler model involving only sending of simple commands initiating genetic programs suggests itself. The following proposal tries to achieve this and involves three basic ideas.

1. The idea of password and addressing is familiar from ordinary computers. Collection of frequencies as password/address allows to reach tuned targets without specific targeting of the command. This is a dramatic improvement to the previous model.
2. Password and fractal addressing realized in terms of frequencies coupling resonantly (already in the original model: I did not however realize the implications of resonant coupling!) and the hierarchy of Planck constants to realize the hierarchical addressing. I have discussed analogous addressing based on information molecules and their receptors at the biochemical level to realize magnetic flux tube connections between sender and target inside organism (hormonal action would be very analogous to what I am proposing here).
3. Becker's DC currents as supra currents flowing along DNA and activated optimally when the incoming laser light has polarization parallel to DNA's local direction, activation of super currents would mean activation of the gene. This is second new element to the original model.

In the following I discuss this with more details.

## 4.1 The analogy with ordinary computer

Consider first the analog of remote mental interactions for ordinary computer. Computer sends a password to the other computer and after that it can use it to run programs of the other computer. Whistling to a dog is another example: extremely simple command activates arbitrary complex programs.

In the recent case electromagnetic radiation with a given frequency coupling resonantly like radio signal to a tuned radio receiver would be the simplest command activating the target. There would no need to specify the direction or distance of the target precisely since essentially mass communications would be in question: intent would be enough. Password could consist of several frequencies which must be received simultaneously by the target before it would activate and tunes to receive more frequencies representing simple commands - perhaps acting on the intronic portion of DNA and activating the genome to remote gene expression or something else such as activating DNAs of other cells by sending similar em addresses!

I have discussed topological quantum computer programs based on braiding could look like in this framework [L3]. Also here addressing but now realized as information molecule-receptor pair would play a key role.

## 4.2 Hierarchy of Planck constants and hierarchical addressing

Fractal hierarchy of frequencies (in Peter's experiment laser light induced generation of radiation at frequencies down to about 10 kHz) would allow to transform passwording to addressing. Very naively, the longest wavelengths: about  $10^4$  meters would reach the tuned receivers in nearly the same phase in a region of this size. One would have some subregions in tune. The shorter wavelengths would allow to pinpoint the tuned receivers inside each of these subregions and so on. This would be fractal addressing with most significant bits correspond to the longest wavelengths. Only those receivers which would be tuned to all frequencies would start to express the gene in the case of AND logic. Of course, also other Boolean functions of tuned-not tuned bits can be considered.

A good guess is that all photons correspond to same energy of visible photon and only Planck constant varies. For ordinary value of Planck constant one would have a photon with wavelength of order size scale of single cell, and the frequencies in this range would select single gene in the genome of particular kind of cell, say neuron within particular region of brain.

In Peter Gariaev's experiment involving 2 eV incoming red laser light the outgoing photons would have same energy but larger Planck constant so that also wavelengths would be longer and range down to at least  $3 \times 10^4$  meters corresponding to radiofrequency scale of 10 kHz. What is interesting that 2 eV is 4 times the nominal value of the metabolic energy quantum of .5 eV identifiable as zero point kinetic energy of electron or proton for the p-adic length scale  $L(151)$  corresponding to cell membrane thickness and Gaussian Mersenne  $M_{151} = (1 + i)^{151} - 1$ . Could it be that 2 eV could be preferred photon energy or is its use simply due to the unavailability of continuous frequency spectrum for laser light. And why the laser light induces the generation of the command inducing remote gene expression?

This picture conforms with Peter's experiment and with the reports of Benveniste and followers about the possibility of representing homeopathic remedy using very low frequency spectrum - presumably cyclotron frequencies - assignable to remedy. These frequencies would be addresses for genes activating genes transcribing building bricks of biomolecules of immune response eliminating the substance from the organism. The proposal could be seen as a generalization of Benveniste's observation and realization of wave DNA proposal.

## 4.3 DNA supra currents and activation of genes by Becker mechanism

The third building brick of the model would be quantum model for Becker currents [L4] as supra currents or quantal DC currents [L4]: also this element is new. Assume - in accordance with the general vision - that these supra currents can flow also along the strands of many-sheeted DNA (flux sheets associated with the strand, entire hierarchy labelled by the values of  $\hbar$ ). Assume also that the interaction of polarized photons addressing for genes with DNA is such that the electric fields of DNA flux tube and "massless extremal" representing laser beam superpose and charges (electrons) experience the superposition of field already present and the field of ME. If the net electric field is near

criticality originally (think as analog neuronal membrane) and becomes over-critical, quantal Becker current starts to flow and the machinery responsible for gene activation is activated.

This means also the activation of metabolic machinery since the acceleration of electrons in the electric field gives them energy making possible a transfer to smaller space-time sheets where they form Cooper pair like states with negentropic entanglement. Metabolic energy corresponds to zero point kinetic energy and negentropic entanglement is relevant from the point of view of consciousness: in the case of healing understood as a regeneration of negentropic resources this aspect is especially important. This mechanism generates high energy phosphate bonds in ATP and the decay  $\text{ATP} \rightarrow \text{ADP}$  liberates the metabolic energy and destroys the negentropic entanglement possibly associated with ATP so that the second law in generalized form [L2] allowing local generation of genuine negentropy (but assigned to information carried by entanglement defining a quantum rule) wins after all.

It could also happen that the decay of ATP generates dark photon or photons absorbed by cyclotron condensate at magnetic flux tube. The excited state is non-local single particle excitation and involves very simple negentropic entanglement between the particles of the condensate. In this case the negentropy of ATP would be transformed to the negentropy of the magnetic flux tube or even several of them if large value of Planck constant is associated with the photon. This mechanism could allow the generation of negentropic entanglement associated with attention. The storage of metabolic energy in photosynthesis could involve similar excitation of cyclotron state at the first step. The most plausible candidate is cyclotron condensate for electron Cooper pairs. Also electrons filling state up to some Fermi energy could be in question. In this case the excitations would be excitation in longitudinal degrees of freedom of the flux tube generating current.

## 5 DNA, speech, music, and ordinary sound

Peter Gariaev's group has made rather dramatic claims about DNA during years [I4, I7, I6, I8].

1. The group has proposed that the statistical distributions of nucleotides and codons in the intronic portion of DNA resemble the distribution of letters and words in the natural languages [I8]. For instance, it is proposed that Zipf law [J7] applying to natural languages applies to the distributions of codons in the intronic portion of DNA. One can study the popularity of the words in natural languages and order them against their popularity. Zipf law states that the integer characterizing popularity is in constant proportion to the number of times it appears in given long enough text.
2. It has been also claimed that DNA can be reprogrammed using modulated laser light or even radio waves. I understand that reprogramming means a modified gene expression. Gariaev's group indeed proposes that the meaning of the third nucleotide (having a rather low significance in the DNA-aminoacid correspondence) in the genetic codon depends on the context giving rise to a context dependent translation to amino-acids. This is certainly a well-known fact for certain variants of the genetic code. This context dependence might make possible the re-programming. The notion of dark DNA allows to consider much more radical possibility based on the transcription of dark DNA to mRNA followed by translation to aminoacids. This could effectively replaced genes with new ones.
3. Also the modulation of the laser light by speech is claimed to have the re-programming effect. The broad band em wave spectrum resulting in the scattering of red laser light on DNA is reported to have rather dramatic biological effects. The long wave length part of this spectrum can be recorded and transformed to sound waves and these sound waves are claimed to have the same biological effects as the light. The proposal is that acoustic solitons propagating along DNA represent this effect on DNA.

I do not have the competence to make statements about the plausibility of these claims. TGD view about quantum biology makes also rather strong claims. The natural question is however whether a justification for the claims of Gariaev and collaborators could be found in TGD framework? In particular, can one say about possible effects of sound on DNA. One intriguing fact about sound perception is that music and speech have meaning whereas generic sounds do not. Could one say something interesting about how this meaning is generated at the level of DNA?

## 5.1 Basic picture

Before continuing it is good to restate the basic TGD inspired ideas about the generation of meaning.

1. The generation of the negentropic entanglement is the correlate for the experience of the meaning. In the model inspired by Becker's findings, the generation of negentropic entanglement involves a generation of supra currents along flux tubes moving in the electric field parallel to them. This is a critical phenomenon taking place when the voltage along the flux tube is near critical value. The generation of nerve pulse near critical value of the resting potential is one example of this criticality. Becker's direct currents involved with the healing of wounds is another example.

The flow of the supra current gives rise to the acceleration of charges along the flux tubes and generation of Cooper pairs or even many-electrons systems at smaller space-time sheets in negentropically entangled state and carrying metabolic energy quantum as zero point kinetic energy. The period of negentropic entanglement gives rise to a conscious experience to which one can assign various attributes such as understanding, attention, and so on. Negentropic entanglement would measure the information contained by a rule having as instances the state pairs in the quantum superposition defining the entangled state. When the period of negentropic entanglement ceases, the metabolic energy is liberated.

2. Remote activation of DNA by analogs of laser beams is another essential piece of TGD inspired quantum biology. In the proposed addressing mechanism a collection of frequencies serves as a password activating intronic portions of DNA. This would take place via a resonance for the proposed interaction between photons and dark supra currents flowing along magnetic flux tubes and perhaps also along DNA strands or flux tubes parallel to them. The superposition of electric fields of photons (massless extremals) with the electric fields parallel to flux tubes (so that massless extremals serving as correlates for laser beams would traverse the flux tube in orthogonal direction).
3. The flux tubes, and more generally flux sheets labelled by the value of Planck constant, and along which the radiation arrives would be transversal to DNA and contain DNA strands. This kind of flux tubes and sheets also define the connections to the magnetic body, and form parts of it. A given flux sheet would naturally select the portion of DNA, which is activated by the radiation: it could be a portion of intronic part of DNA activating in turn a gene. These flux tubes and sheets could be connected to the lipids of nuclear and cell membranes - also cell membranes of other cells - as assumed in the model of DNA as topological quantum computer [K2]. The sheets could also give rise to a hierarchy of genomes - besides genome one would have super-genome in which genomes are organelles are integrated by flux sheets to a large coherently expressed structure containing individual genomes like page of a book contains lines of text. These pages would be in turn organized to a book - hyper-genome as I called it. One could have also libraries, etc... There would fractal flux quanta inside flux quanta structure.

## 5.2 Phonons and photons In TGD Universe

Consider next phonons and their coupling to photons in TGD Universe.

1. Sound waves could quite well transform to electromagnetic radiation since living matter is piezo-crystal transforming sound to radiation and vice versa. Microwave hearing represents an example of this kind transformation. This would require that photons of given energy and varying value of Planck constant couple to phonons with the same energy, Planck constant, and frequency.
2. Whether one can assign to phonons a non-standard value of Planck constant is not quite clear, but there seems to be no reason preventing this. If so, even photons of audible sounds would have energies above thermal threshold and have direct quantal effects on living matter if they have same Planck constant as the photons with same frequency.
3. Acoustic phonons represent longitudinal waves and this would require longitudinal photons. In Maxwell's electrodynamics they are not possible but in TGD framework photon is predicted to have a small mass and also longitudinal photons are possible.

4. For general condensed matter systems one can have also optical phonons for which the polarization is orthogonal to the wave vector and these could couple to ordinary photons. The motion of the charged particles in the electromagnetic field of massless extremal (topological light ray) would be a situation in which phonons and photons accompany each other. This would make possible the piezo-electric mechanism.

Under these assumptions the collections of audible frequencies could also represent passwords activating the intronic portion of the genome and lead to gene expression or some other activities. If one believes on the hypothesis that DNA acts like topological quantum computer based on the braid strand connections between nucleotides in the intronic portion of DNA with the lipids of the nuclear and/or cell membranes, also topological quantum computation type processes could be activated by the collections of sound frequencies [K2].

### 5.3 What distinguishes speech and music from sounds without meaning?

Speech and music are very special form of sound in that they have direct meaning. The more one thinks about these facts, the more non-trivial they look. For music - say singing - the frequency of the carrier wave is piecewise constant whereas for speech it remains constant and the amplitude modulation is important. In fact, by slowing down the recorded speech, one gets the impression that carrier frequency is actually modulated like in chirp (frequency goes down and covers a range of frequencies). What is the mechanism giving to speech and music its meaning and in this manner distinguishes them from other sounds?

Besides the frequency also phase is important for both speech and music experience. Speech and reverse speech sound quite different the intensity in frequency space is same. Therefore the relative phases associated with the Fourier coefficients of various frequencies must be important. For music simple rational multiples of the fundamental define the scale. Could it be that also the frequencies relevant to the comprehension of speech correspond to these rational multiples?

Suppose that one indeed believes on the proposed vision based on the fundamental role of negentropic entanglement in generation of meaning and takes seriously the proposed mechanisms for generating it. Can one understand why music and speech differ from general sounds and what distinguishes between them?

1. With these assumptions suitable collections of frequencies sound wave would indeed activate the intronic portion of DNA by generating negentropic entanglement. Also other dark flux tubes than those assignable to DNA are involved. For instance, hair cells responsible for hearing of sounds around particular frequencies could involve flux tubes and utilize similar mechanism. Allowing only hair cells would define the conservative option. On the other hand, one could well claim that what happens in ear has nothing to do with the understanding of the speech and music, it could take place only at the level of neuronal nuclei.
2. Could the direct interaction of sound waves with magnetic flux tubes generate the experiences of speech and music? In other words, assign meaning to sounds? The criterion for sound to have an interpretation as speech or music would be that it contains the resonance frequencies needed to activate the DNA, or more generally generate dark super currents generating Cooper pairs in this manner loading metabolic energy storages. This would apply to both speech and musical sounds.
3. The pitch of the speech and musical sound can vary. We are aware of the key of the music piece and of modulations of the key and remember the starting key, and it is highly satisfactory to make a return to "home" defined by the original key. This would imply that the overall scale of the collection of frequencies can be varied and that the pitch of the speech defines a natural expectation value of this scale. For persons possessing so called absolute ear this scaling symmetry would be broken in a well-defined sense.
4. Musical scales involve frequencies coming as rational multiples of the basic frequency. Octaves - power of two multiples - of the frequency can be said to be equivalent as far as musical experience is considered. One might understand the special role of rational multiples of the basic frequency if the Fourier components have same phase periodically so that the experience is invariant

under discrete time translations. This requires commensurable frequencies expressible as rational multiples of the same fundamental frequency. The preferred role of p-adic primes coming as powers of two could relate to the octave phenomenon.

5. Are the relative phases of different Fourier components important for music experience? If one requires a periodical occurrence of maximal possible intensity (maximal constructive interference) then the relative phases must vanish at the values of time for maximal possible intensity. What seems essential that the presence of commensurate frequencies gives rise to time translation invariant sensation whereas speech consists of pulses.

#### 5.4 Are speech and music quantum duals like position and momentum?

Frequencies are crucial for music experience. In the case of speech the relative phases are very important as the example of reverse speech demonstrates. How a given phoneme is heard is determined to high degree by the frequency spectrum in the beginning of the phoneme (this distinguishes between consonants). Vowels are nearer to notes in vocalization. Speech consists of pulses and destructive interference between different frequencies is required to generate pulses and different pulse shapes so that phase information is important. At least the harmonics of the basic rational multiples of the fundamental are necessary for speech.

One can criticize the previous discussion in that it has been completely classical. Phase and frequency are in wave mechanics canonically conjugate variables analogous to position and momentum. Is it really possible to understand the difference between music and speech purely classically by assuming that one can assign to sound waves both frequencies and phases simultaneously - just like one assigns to a particle sharp values of both momentum and position? Or should one use either representation either in terms numbers of phonons in different modes labelled by frequencies or as coherent states of phonons with ill defined phonon numbers but well defined amplitudes? Could the coherent states serve as the analogs of classical sound waves. Speech would be as near as possible to classical sound and music would be quantal. Of course, there is a large variety of alternative choices of basis states between these two extremes as a specialist in quantum optics could tell.

Suppose that this picture is more or less correct. What could be the minimal scenario allowing to understand the differences between speech and music?

1. Only a subset of frequencies could activate DNA (or if one wants to be conservative, the hair cells) also in the case of speech. One could still pick up important frequencies for which the ratios are simple rational numbers as in the case of musical scale plus their harmonics. If this assumption is correct, then speech from which all frequencies except for the harmonics of the simple rational multiples of the fundamental are removed, should be still be comprehensible as speech. The pitch of the speech would determine a good candidate for the fundamental frequency.
2. The harmonics of frequencies activating DNA would be crucial for speech. Harmonics are present also in music and their distribution allows to distinguish between different instruments and persons. The deviation of musical notes from ideal Fock states would correspond to this.
3. The naive guess is that the simple rational multiples of fundamental and the possibility of having their harmonics could be reflected in the structure of intronic portions of DNA as repetitive structures of various sizes. This cannot be the case since the wavelengths of ordinary photons would be so small that the energies would be in keV range. Neither is this expected to be the case. It is magnetic flux tubes and sheets traversing the DNA which carry the radiation and the natural lengths assignable to these flux quanta should correspond to the wave lengths. The larger, the flux quantum, the lower the frequency and the larger the value of Planck constant. Harmonics of the fundamental would appear for given flux tube length naturally.

The DNA strands and flux tubes and sheets form a kind of electromagnetic music instrument with flux quanta taking the role of guitar strings and DNA strands and other structures such as lipids and possible other molecules to which flux tubes get attached taking the role of frets in guitar. This analogy suggests that for wave lengths measured in micrometers the basic frequencies correspond to the distances between "frets" defined by cell and nuclear membranes

in the tissue in the scale of organism. This would relate the spectrum of resonance frequencies to the spectrum of distances between DNAs in the tissue.

For wave lengths corresponding to very large values of Planck constant giving rise to frequencies in VLF and ELF range and corresponding also to audible frequencies, the preferred wave lengths would correspond to lengths of flux quanta in Earth size scale. One should understand whether the quantization of these lengths in simple rational ratios could take place for the preferred extremals.

4. Could the pulse shape associated with massless extremals (MEs, topological light rays) allow to distinguish classically between speech and music at the level of space-time correlates? Linear superposition of Fourier components in the direction of ME is possible and this allows to speak about pulse shape. It allows also the notions of coherent state and Fock state for given direction of wave vector. Essential would be the restriction of the superposition of fields in single direction of propagation to be distinguished from the superposition of the effects of fields associated with different space-time sheets on multiply topologically condensed particle. Maybe this would allow to make testable predictions.

## REFERENCES

### Theoretical Physics

- [B1] Airy functions. [http://en.wikipedia.org/wiki/Airy\\_function](http://en.wikipedia.org/wiki/Airy_function).

### Biology

- [I1] Dr. Phil Callahan on Power of Paramagnetism. *Nexus*. <http://www.nexusmagazine.com>, February 2003.
- [I2] Acupuncture's Molecular Effects Pinned Down: New Insights Spur Effort to Boost Treatment's Impact Significantly. Science News. <http://www.sciencedaily.com/releases/2010/05/100530144021.htm>, 2010.
- [I3] N. Goldman et al. Adenosine A1 receptors mediate local anti-nociceptive effects of acupuncture. *Nature Neuroscience*. <http://www.nature.com/neuro/journal/v13/n7/full/nn.2562.html>, 13:883–888, 2010.
- [I4] P. Gariaev et al. *The DNA-wave biocomputer*, volume 10. CHAOS, 2001.
- [I5] T. I. Karu. *The Science of Low-Power Laser Therapy*. Gordon and Breach, Sci. Publ. <http://www.laserhealthsystems.com/Dr.%20Tiina%20Karu%20Presentation.htm>, London, 1998.
- [I6] A.V. Tovmash P. P. Gariaev, G. G. Tertishni. Experimental investigation in vitro of holographic mapping and holographic transposition of DNA in conjunction with the information pool encircling DNA. *New Medical Tehcnologies*, 9:42–53, 2007.
- [I7] G. G. Komissarov A. A. Berezin A. A. Vasiliev P. P. Gariaev, V. I. Chudin. Holographic Associative Memory of Biological Systems. *Proceedings SPIE - The International Society for Optical Engineering. Optical Memory and Neural Networks*, pages 280–291, 1991.
- [I8] M. U. Maslow P. P. Gariaev. Fractal Presentation of Natural Language Texts and Genetic Code, 2nd International Conference on Quantitative Linguistics. *QUALICO '94*, pages 193–194, 1994.

### Neuroscience and Consciousness

- [J1] Pflanzenwachstum durch Elektrofild. <http://www.s-line.de/homepages/keppler/elektrofild.htm>.



- [J2] R. O. Becker and G. Selden. *The Body Electric: Electromagnetism and the Foundation of Life*. William Morrow & Company, Inc., New York, 1990.
- [J3] M. L. Recce J. O'Keefe. Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus*. <http://www3.interscience.wiley.com/cgi-bin/abstract/109711451/ABSTRACT>, (3):317–330, 2004.
- [J4] L. Sidorov and K. W. Chen. Biophysical Mechanisms of Genetic Regulation: Is There a Link to Mind-Body Healing? *DNA Decipher Journal*, 2(2):177–205, 2012.
- [J5] Y. Yamaguchi. Laboratory for dynamics of emergent intelligence. [http://www.brain.riken.jp/reports/annual\\_reports/2003/2003-doc/0243.pdf](http://www.brain.riken.jp/reports/annual_reports/2003/2003-doc/0243.pdf), 2003.
- [J6] Y. Yamaguchi. Neural dynamics in Hippocampal Memory Formation via Theta Rhythm. <http://www.brain.riken.jp/bsi-news/bsinews22/no22/speciale.html>, 2003.
- [J7] G. K. Zipf. *Psycho-Biology of Languages*. MIT Press, 1965.

## Books related to TGD

- [K1] M. Pitkänen. About the New Physics Behind Qualia. In *Quantum Hardware of Living Matter*. Onlinebook. [http://tgdtheory.com/public\\_html/bioware/bioware.html#newphys](http://tgdtheory.com/public_html/bioware/bioware.html#newphys), 2006.
- [K2] M. Pitkänen. DNA as Topological Quantum Computer. In *Genes and Memes*. Onlinebook. [http://tgdtheory.com/public\\_html/genememe/genememe.html#dnatqc](http://tgdtheory.com/public_html/genememe/genememe.html#dnatqc), 2006.
- [K3] M. Pitkänen. Genes and Memes. In *Genes and Memes*. Onlinebook. [http://tgdtheory.com/public\\_html/genememe/genememe.html#genememec](http://tgdtheory.com/public_html/genememe/genememe.html#genememec), 2006.
- [K4] M. Pitkänen. Homeopathy in Many-Sheeted Space-Time. In *Bio-Systems as Conscious Holograms*. Onlinebook. [http://tgdtheory.com/public\\_html/hologram/hologram.html#homeoc](http://tgdtheory.com/public_html/hologram/hologram.html#homeoc), 2006.
- [K5] M. Pitkänen. Nuclear String Hypothesis. In *p-Adic Length Scale Hypothesis and Dark Matter Hierarchy*. Onlinebook. [http://tgdtheory.com/public\\_html/paddark/paddark.html#nuclstring](http://tgdtheory.com/public_html/paddark/paddark.html#nuclstring), 2006.
- [K6] M. Pitkänen. Quantum Model for Nerve Pulse. In *TGD and EEG*. Onlinebook. [http://tgdtheory.com/public\\_html//tgdeeg/tgdeeg/tgdeeg.html#pulse](http://tgdtheory.com/public_html//tgdeeg/tgdeeg/tgdeeg.html#pulse), 2006.
- [K7] M. Pitkänen. *TGD Based View About Living Matter and Remote Mental Interactions*. Onlinebook. [http://tgdtheory.com/public\\_html/tgdlian/tgdlian.html](http://tgdtheory.com/public_html/tgdlian/tgdlian.html), 2012.
- [K8] M. Pitkänen and P. Gariaev. Quantum Model for Remote Replication. In *Genes and Memes*. Onlinebook. [http://tgdtheory.com/public\\_html/genememe/genememe.html#remotereplication](http://tgdtheory.com/public_html/genememe/genememe.html#remotereplication), 2011.

## Articles about TGD

- [L1] M. Pitkänen. DNA Waves and Water. [http://tgdtheory.com/public\\_html/articles/mont.pdf](http://tgdtheory.com/public_html/articles/mont.pdf), 2011.
- [L2] M. Pitkänen. Generalization of thermodynamics allowing negentropic entanglement and a model for conscious information processing. <http://tgdtheory.com/articles/thermolife.pdf>, 2011.
- [L3] M. Pitkänen. How to build a quantum computer from magnetic flux tubes? . <http://tgdtheory.com/articles/fluxtubecomputer.pdf>, 2012.
- [L4] M. Pitkänen. Quantum Model for the Direct Currents of Becker. [http://tgdtheory.com/public\\_html/articles/DCbio.pdf](http://tgdtheory.com/public_html/articles/DCbio.pdf), 2012.