

Official top quark and toponium as particles of M_{89} hadron physics?

March 13, 2026

Matti Pitkänen

orcid:0000-0002-8051-4364.

email: matpitka6@gmail.com,

url: http://tgdtheory.com/public_html/,

address: Valtatie 8, as. 2, 03600, Karkkila, Finland.

Contents

1	Introduction	1
2	Toponium anomaly as an indication for M_{89} hadron physics	2
2.1	What toponium is?	2
2.2	TGD predicts a fractal hierarchy of standard model physics	3
2.3	Could the official top quark be the u quark of M_{89} hadron physics?	3

Abstract

The anomalies of the standard model have begun to accumulate. The existence of toponium is the newest anomaly. The lifetime of the official top quark candidate is too short for the toponium to form by hadronic interactions. The new view of quark color predicted by TGD, predicts an entire fractal hierarchy of hadron physics characterized by the color multiplets defined by color partial waves in CP_2 . p-Adic length scale hypothesis, having now number theoretic justification, suggests that ordinary and Gaussian Mersenne primes label these copies characterized by a p-adic length scale coming as some powers of 2. M_{89} hadron physics corresponds to a hadronic mass scale, which is 512 times that for ordinary hadron physics. For top quark there exist unorthodox candidates at masses 28 and 30 GeV and there is also a candidate for toponium with mass 56 GeV. The official top quark has suspiciously large mass: could it correspond to u quark of M_{89} hadron physics and could toponium be one state of isospin triplet representing pion or ρ meson of M_{89} hadron physics.

1 Introduction

I watched an excellent video about what we have learned at LHC (see). Three runs RUN1, RUN2, and RUN3 have been completed and now we know where the limits for the applicability of the Standard Model are.

The immediate successor of LHC will be high-luminosity LHC operating from 2029- 2030 onwards for ten years. Future circular collider (FCC) will start to operate in the late 2040s. Electrons and positrons will collide and the collider (Higgs factory) will act as a high precision collider.

The philosophy is that high precision might allow us to develop a theory allowing us to solve the various anomalies of the standard model. In future, the experimentalists would not be merely

testing whether a given extension of the standard model might solve some anomalies but trying to identify more general deviations from the standard model. But is this enough? What has been lacking from theoretical physics since the times of Einstein and his contemporaries, is philosophical thinking challenging the basic assumptions. Can one make progress by merely measuring more precisely?

The video explains the basic anomalies. The anomalies are also discussed in detail by Crivellin and Mellado [C2]. The following list defines the boundaries of the region of phenomena that the standard model can explain.

1. Toponium exists although it should not.
2. W mass deviates from the predicted mass.
3. $g-2$ anomaly of muon is claimed to disappear in lattice calculations using only quarks and gluons but does not disappear when hadronic data are used as an input.
4. Lepton universality is violated in some meson decays.
5. Penta and tetra quarks, whose existence is not denied but not predicted by the standard model.
6. There are anomalies associated with the CP violation of the CKM matrix.
7. The axions, proposed to solve the problem due to the strong CP violation predicted by QCD, have not been found and the strong CP violation is too weak to explain matter antimatter asymmetry.
8. Quark-gluon plasma predicted by QCD did not behave like gas but a perfect liquid and the transition to quark gluon plasma seems to occur at several energies rather than single phase transition point.
9. SUSY was believed to solve the hierarchy problem involving the fine tuning of the Higg couplings but no evidence for SUSY particles was found.
10. WIMPs as candidates for galactic dark matter have not been found.

2 Toponium anomaly as an indication for M_{89} hadron physics

I have discussed various anomalies from the TGD point of view in various articles. Here I will consider only the discovery of the toponium, which is one of the latest surprises.

2.1 What toponium is?

The Standard model does not deny toponium's existence but according to the standard intuition it should not exist.

1. The lifetime of the top quark is too short for the formation of toponium. There are of course proposals for solving this and also other anomalies but the problem is that these proposals typically solve only one anomaly. The lifetime of the standard top quark candidate with mass $m \simeq 172.5$ GeV is $\tau = 5 \times 10^{-25}$ s. This time is shorter than required for QCD hadronization processes ($10^{-23} - 10^{-24}$ s). This is why it has been believed that toponium does not exist.
2. The toponium was however discovered both by LHC and ATLAS and its lifetime is estimated to be 2.5×10^{-25} s. Toponium is suggested to be a quasi-bound state or a resonance appearing when top quarks are produced very near to the threshold energy (see this and this). Toponium decay is triggered by a weak decay of one of its constituents rather than being a strong decay. Both ATLAS and CMS verified the existence of this state with a resonance width of about 3 GeV.

2.2 TGD predicts a fractal hierarchy of standard model physics

Consider now the basic ideas of TGD view of hadron physics and standard model in general. TGD leads to almost inescapable conclusion that there must exist an entire hierarchy of standard model physics assignable to the triality ± 1 color representations defined by color partial waves of quarks and antiquarks in CP_2 . Leptons would appear in triality 0 color partial waves [L4, ?].

1. The color multiplets of quarks of a given standard model physics would combine to form color triplets, which would serve as building bricks of hadrons of a given hadron physics [L5, L8, L3, L7]. These hadrons would correspond to a hierarchy of p-adic mass scales, proposed to be labelled by ordinary and Gaussian Mersenne primes. The longer the p-adic scale, the higher the dimension of the color multiplet.

For the observed leptons, color representations would combine to form color singlets but also analogs of mesons as bound states of colored leptons might be possible [K5]. Only at energies near CP_2 mass would color deconfinement for incoming and outgoing states be possible.

2. Ordinary hadrons would correspond to the Mersenne prime M_{107} . The nucleon of M_{89} hadron physics would correspond to the mass scale $512m_n$ and therefore to the LHC energy scale. The transition from M_{107} hadron physics to M_{89} hadron physics would take place at quantum criticality. The phase transition usually interpreted as a creation of the quark-gluon phase could correspond to this phase transition [L6]. At quantum criticality the value h_{eff}/h would scale up the Compton length scale of M_{89} hadrons. This would reflect long range quantum fluctuations. This re-interpretation of what has been identified as quark gluon-plasma would solve various anomalies associated with this identification mentioned already in the list of anomalies [L6, L5]. The existence of M_{89} hadron physics can have dramatic implications. For instance, a dramatic modification of the model of the Sun [L2] can be considered.
3. The ratio of the p-adic length scales associated with M_{107} and M_{89} , characterizing the Compton lengths and also defining the geometric size of nucleons as 3-surfaces, is 512. The assumption is that the geometric size of the M_{89} hadron with a large h_{eff} is the same as for M_{107} hadron at quantum criticality implies $h_{eff}/h = 512$. The sizes of M_{89} hadrons would be the same as for ordinary hadrons at quantum criticality for the transition from M_{89} hadron physics to M_{107} hadron physics.
4. I have proposed the identification of various bumps observed at LHC, originally identified first as candidates for SUSY particles but then rejected, in terms of M_{89} mesons [K2, K3].

2.3 Could the official top quark be the u quark of M_{89} hadron physics?

The large mass of the official top quark raises the question whether it could be M_{89} quark created at quantum criticality.

1. A natural guess is that the lifetime of top quark at quantum criticality is scaled up $h_{eff}/h = 512$ to $.25 \times 10^{-21}$ s. The corresponding distance scale would be $.75 \times 10^{-13}$ m, which is longer than the nuclear size scale!
2. A reasonable guess is that the hadronization time scale for M_{89} is for h_{eff}/h scaled down by factor $1/512$ due to decrease of the p-adic length scales. This p-adic length scale corresponds to the geometric size scale of the causal diamond $CD = cd \times CP_2$ assignable to the region in which the phase transition occurs. This local phase transition is discussed in [L8]. The increase $h_{eff} \rightarrow 512h_{eff}$ keeps the geometric time scale associated for hadronization the same as it would be for ordinary hadrons and determined by the p-adic time scale $L(107)$ assignable to ordinary hadrons.

What happens to the rate of hadronization? The phase transition increasing the value of h_{eff} guarantees that the TGD counterpart of perturbative theory, still applies. "Mother Nature loves her theoreticians" [L1] is one way to express this principle. Since the zeroth order term

in the TGD counterpart of the perturbative expansion, giving the classical approximation, does not depend on h_{eff} , the classical approximation improves as h_{eff} increases.

The rate for M_{89} hadronization is proportional to the hadronic mass scale $m(89) = 512m(107)$. Since the geometric time scale is $L(107)$ by quantum criticality, the short lifetime of top does not prevent the formation of toponium. Quantum criticality could quite generally increase the probabilities for the formation of bound states of very short-lived particles.

The basic objection is that the official top quark as M_{89} quark would most naturally correspond to genus $g = 0$ for the partonic 2-surface and serve as a counterpart of u quark. The actual $g = 2$ U type quark should have a lower mass.

1. There is indeed evidence for a top quark-like state at much lower mass from Aleph. The mass is estimated to be about 30 GeV or 28 GeV [C1, ?]. This has motivated the question whether the two candidates for the top quark could correspond to a scaled variant of the top. In the TGD framework, the p-adic length scale hypothesis might allow this [K2] [L3].
2. What about the toponium in this case? There is an old anomaly reported by Aleph at 56 GeV (see <https://arxiv.org/pdf/hep-ph/9608264.pdf>) and there is reference to an old paper: ALEPH Collaboration, D. Buskulic *et al*, CERN preprint PPE/96-052. What was observed was 4-jet events consisting of dijets with invariant mass around 55 GeV. What makes this interesting is that the mass of 28 GeV particle candidates would be one half of the mass of a particle with a mass of 56 GeV particle, quite near to 55 GeV. Could this state be the toponium as $g = 2$ U quark [K2] [L3]?

If this picture is correct, the official top quark would more naturally correspond to the genus $g = 0$ and therefore to M_{89} u quark. Could the poor understanding of the family replication phenomenon and of the origin of the CKM mixing explain this mis-interpretation?

1. The CKM matrix V is empirically determined from charged currents (W decays). The matrix elements of type V_D^U , $U \in \{u, c\}$, $V \in \{d, s\}$ reflect the CKM mixing of d and s quarks. Unitary conditions bring in the matrix elements V_D^b and dependence on top quark mass. Both beta decays and kaon decays provide information about V_D^U , $U \in \{u, c\}$, $V \in \{d, s\}$. These two kinds of constraints lead to slightly different outcomes [C2] for V .
2. Could a wrong identification of the top quark mass cause the discrepancy? In TGD, the official top as the $g = 0$ quark of "dark" M_{89} hadron physics created in the transition to quark-gluon plasma would induce a leakage of probability inducing a genuine violation of the unitarity for the CKM matrix.
3. In TGD, the description of family replication has topological explanation and CKM mixing reduces to topological mixing discussed in [K1, K4]. A model for the transition between M_{107} and M_{89} is needed to see whether the new interpretation can be consistent with what is known about creation of official top quarks.
4. The prediction is that the official top as an M_{89} u quark is accompanied by an M_{89} d quark so that toponium should be a member of an isospin triplet. The M_{89} counterparts of π and ρ mesons should exist. The discovery of the M_{89} d quark, perhaps through the discovery of an isospin triplet for toponium, with nearly the same mass as the official top quark, would force us to take the TGD view seriously.

REFERENCES

Particle and Nuclear Physics

- [C1] Heisner A. Observation of an excess at 30 gev in the opposite sign di-muon spectra of $z \rightarrow b\bar{b} + x$ events recorded by the aleph experiment at lep, 2016. Available at: <https://tinyurl.com/hy8ugf4>.

- [C2] Mellado B. Crivellin A. Anomalies in Particle Physics, 2025. Available at: <https://arxiv.org/abs/2309.03870>.

Books related to TGD

- [K1] Pitkänen M. Construction of elementary particle vacuum functionals. In *p-Adic Physics*. <https://tgdtheory.fi/tgdhtml/Bpadphys.html>. Available at: <https://tgdtheory.fi/pdfpool/elvafu.pdf>, 2023.
- [K2] Pitkänen M. New Physics Predicted by TGD: Part I. In *p-Adic Physics*. <https://tgdtheory.fi/tgdhtml/Bpadphys.html>. Available at: <https://tgdtheory.fi/pdfpool/TGDnewphys1.pdf>, 2023.
- [K3] Pitkänen M. New Physics Predicted by TGD: Part II. In *p-Adic Physics*. <https://tgdtheory.fi/tgdhtml/Bpadphys.html>. Available at: <https://tgdtheory.fi/pdfpool/TGDnewphys2.pdf>, 2023.
- [K4] Pitkänen M. p-Adic Particle Massivation: Hadron Masses. In *p-Adic Physics*. <https://tgdtheory.fi/tgdhtml/Bpadphys.html>. Available at: <https://tgdtheory.fi/pdfpool/mass3.pdf>, 2023.
- [K5] Pitkänen M. The Recent Status of Lepto-hadron Hypothesis. In *p-Adic Physics*. <https://tgdtheory.fi/tgdhtml/Bpadphys.html>. Available at: <https://tgdtheory.fi/pdfpool/leptc.pdf>, 2023.

Articles about TGD

- [L1] Pitkänen M. About the TGD based views of family replication phenomenon and color confinement. https://tgdtheory.fi/public_html/articles/emuanomaly.pdf, 2023.
- [L2] Pitkänen M. Some solar mysteries. https://tgdtheory.fi/public_html/articles/Haramein.pdf, 2024.
- [L3] Pitkänen M. A refined view of the phenomenology of hadron physics and p-adic mass calculations. https://tgdtheory.fi/public_html/articles/padmass2025.pdf, 2025.
- [L4] Pitkänen M. About the structure of Dirac propagator in TGD. https://tgdtheory.fi/public_html/articles/dirprop.pdf, 2025.
- [L5] Pitkänen M. Comparing the S-matrix descriptions of fundamental interactions provided by standard model and TGD. https://tgdtheory.fi/public_html/articles/hadroQCDTGD.pdf, 2025.
- [L6] Pitkänen M. The findings of RHIC about quark gluon plasma from the TGD point of view. https://tgdtheory.fi/public_html/articles/RHICQGP.pdf, 2025.
- [L7] Pitkänen M. Does the notion of Teichmüller element cure the problem of p-adic mass calculations due to the slight failure of Lorentz invariance? https://tgdtheory.fi/public_html/articles/padmass2026.pdf, 2026.
- [L8] Pitkänen M. TGD counterpart of Feynman diagrammatics applied to QFT limit and CP violation. 2026. Available at: https://tgdtheory.fi/public_html/articles/QFTlimit.pdf.