

Introduction to "Quantum TGD as Infinite-Dimensional Spinor Geometry"

M. Pitkänen,

November 30, 2016

Email: matpitka@luukku.com.

http://tgdtheory.com/public_html/.

Recent postal address: Karkinkatu 3 I 3, 00360, Karkkila, Finland.

Contents

1	Basic Ideas Of Topological Geometrodynamics (TGD)	3
1.1	Basic Vision Very Briefly	3
1.2	Two Vision About TGD And Their Fusion	6
1.2.1	<i>TGD as a Poincare invariant theory of gravitation</i>	6
1.2.2	<i>TGD as a generalization of the hadronic string model</i>	6
1.2.3	<i>Fusion of the two approaches via a generalization of the space-time concept</i>	7
1.3	Basic Objections	8
1.3.1	<i>Topological field quantization</i>	8
1.4	P-Adic Variants Of Space-Time Surfaces	8
1.5	The Threads In The Development Of Quantum TGD	9
1.5.1	<i>Quantum TGD as spinor geometry of World of Classical Worlds</i>	10
1.5.2	<i>TGD as a generalized number theory</i>	12
1.6	Hierarchy Of Planck Constants And Dark Matter Hierarchy	16
1.6.1	<i>Dark matter as large \hbar phases</i>	16
1.6.2	<i>Hierarchy of Planck constants from the anomalies of neuroscience and biology</i>	16
1.6.3	<i>Does the hierarchy of Planck constants reduce to the vacuum degeneracy of Kähler action?</i>	17
1.6.4	<i>Dark matter as a source of long ranged weak and color fields</i>	18
1.7	Twistors And TGD	18
2	Bird's Eye Of View About The Topics Of The Book	19

3 Sources	20
4 The contents of the book	20
4.1 Identification of the Configuration Space Kähler Function	20
4.2 About Identification of the Preferred extremals of Kähler Action	21
4.3 Construction of WCW Kähler Geometry from Symmetry Principles	21
4.4 WCW Spinor Structure	22
4.5 Recent View about Kähler Geometry and Spin Structure of WCW	26
4.6 The Classical Part of the Twistor Story	26
4.7 Unified Number Theoretical Vision	27
4.8 Knots and TGD	27

1 Basic Ideas Of Topological Geometrodynamics (TGD)

Standard model describes rather successfully both electroweak and strong interactions but sees them as totally separate and contains a large number of parameters which it is not able to predict. For about four decades ago unified theories known as Grand Unified Theories (GUTs) trying to understand electroweak interactions and strong interactions as aspects of the same fundamental gauge interaction assignable to a larger symmetry group emerged. Later superstring models trying to unify even gravitation and strong and weak interactions emerged. The shortcomings of both GUTs and superstring models are now well-known. If TGD - whose basic idea emerged 37 years ago - would emerge now it would be seen as an attempt trying to solve the difficulties of these approaches to unification.

The basic physical picture behind TGD corresponds to a fusion of two rather disparate approaches: namely TGD as a Poincare invariant theory of gravitation and TGD as a generalization of the old-fashioned string model. The CMAP files at my homepage provide an overview about ideas and evolution of TGD and make easier to understand what TGD and its applications are about (<http://tgdtheory.fi/cmaphtml.html> [L1]).

1.1 Basic Vision Very Briefly

T(opological) G(eometro)D(ynamics) is one of the many attempts to find a unified description of basic interactions. The development of the basic ideas of TGD to a relatively stable form took time of about half decade [K1].

The basic vision and its relationship to existing theories is now rather well understood.

1. Space-times are representable as 4-surfaces in the 8-dimensional imbedding space $H = M^4 \times CP_2$, where M^4 is 4-dimensional (4-D) Minkowski space and CP_2 is 4-D complex projective space (see Appendix).
2. Induction procedure (a standard procedure in fiber bundle theory, see Appendix) allows to geometrize various fields. Space-time metric characterizing gravitational fields corresponds to the induced metric obtained by projecting the metric tensor of H to the space-time surface. Electroweak gauge potentials are identified as projections of the components of CP_2 spinor connection to the space-time surface, and color gauge potentials as projections of CP_2 Killing vector fields representing color symmetries. Also spinor structure can be induced: induced spinor gamma matrices are projections of gamma matrices of H and induced spinor fields just H spinor fields restricted to space-time surface. Spinor connection is also projected. The interpretation is that distances are measured in imbedding space metric and parallel translation using spinor connection of imbedding space.

The induction procedure applies to octonionic structure and the conjecture is that for preferred extremals the induced octonionic structure is quaternionic: again one just projects the octonion units. I have proposed that one can lift space-time surfaces in H to the Cartesian product of the twistor spaces of M^4 and CP_2 , which are the only 4-manifolds allowing twistor space with Kähler structure. Now the twistor structure would be induced in some sense, and should co-incide with that associated with the induced metric. Clearly, the 2-spheres defining the fibers of twistor spaces of M^4 and CP_2 must allow identification: this 2-sphere defines the S^2 fiber of the twistor space of space-time surface. This poses constraint on the imbedding of the twistor space of space-time surfaces as sub-manifold in the Cartesian product of twistor spaces.

3. Geometrization of quantum numbers is achieved. The isometry group of the geometry of CP_2 codes for the color gauge symmetries of strong interactions. Vierbein group codes for electroweak symmetries, and explains their breaking in terms of CP_2 geometry so that standard model gauge group results. There are also important deviations from standard model: color quantum numbers are not spin-like but analogous to orbital angular momentum: this difference is expected to be seen only in CP_2 scale. In contrast to GUTs, quark and lepton numbers are separately conserved and family replication has a topological explanation in terms of topology of the partonic 2-surface carrying fermionic quantum numbers.

M^4 and CP_2 are unique choices for many other reasons. For instance, they are the unique 4-D space-times allowing twistor space with Kähler structure. M^4 light-cone boundary allows a huge extension of 2-D conformal symmetries. Imbedding space H has a number theoretic interpretation as 8-D space allowing octonionic tangent space structure. M^4 and CP_2 allow quaternionic structures. Therefore standard model symmetries have number theoretic meaning.

4. Induced gauge potentials are expressible in terms of imbedding space coordinates and their gradients and general coordinate invariance implies that there are only 4 field like variables locally. Situation is thus extremely simple mathematically. The objection is that one loses linear superposition of fields. The resolution of the problem comes from the generalization of the concepts of particle and space-time.

Space-time surfaces can be also particle like having thus finite size. In particular, space-time regions with Euclidian signature of the induced metric (temporal and spatial dimensions in the same role) emerge and have interpretation as lines of generalized Feynman diagrams. Particle in space-time can be identified as a topological inhomogeneity in background space-time surface which looks like the space-time of general relativity in long length scales.

One ends up with a generalization of space-time surface to many-sheeted space-time with space-time sheets having extremely small distance of about 10^4 Planck lengths (CP_2 size). As one adds a particle to this kind of structure, it touches various space-time sheets and thus interacts with the associated classical fields. Their effects superpose linearly in good approximation and linear superposition of fields is replaced with that for their effects.

This resolves the basic objection. It also leads to the understanding of how the space-time of general relativity and quantum field theories emerges from TGD space-time as effective space-time when the sheets of many-sheeted space-time are lumped together to form a region of Minkowski space with metric replaced with a metric identified as the sum of empty Minkowski metric and deviations of the metrics of sheets from empty Minkowski metric. Gauge potentials are identified as sums of the induced gauge potentials. TGD is therefore a microscopic theory from which standard model and general relativity follow as a topological simplification however forcing to increase dramatically the number of fundamental field variables.

5. A further objection is that classical weak fields identified as induced gauge fields are long ranged and should cause large parity breaking effects due to weak interactions. These effects are indeed observed but only in living matter. A possible resolution of problem is implied by the condition that the modes of the induced spinor fields have well-defined electromagnetic charge. This forces their localization to 2-D string world sheets in the generic case having vanishing weak gauge fields so that parity breaking effects emerge just as they do in standard model. Also string model like picture emerges from TGD and one ends up with a rather concrete view about generalized Feynman diagrammatics. A possible objection is that the Kähler-Dirac gamma matrices do not define an integrable distribution of 2-planes defining string world sheet.

An even stronger condition would be that the induced classical gauge fields at string world sheet vanish: this condition is allowed by the topological description of particles. The CP_2 projection of string world sheet would be 1-dimensional. Also the number theoretical condition that octonionic and ordinary spinor structures are equivalent guaranteeing that fermionic dynamics is associative leads to the vanishing of induced gauge fields.

The natural action would be given by string world sheet area, which is present only in the space-time regions with Minkowskian signature. Gravitational constant would be present as a fundamental constant in string action and the ratio $\hbar/G/R^2$ would be determined by quantum criticality condition. The hierarchy of Planck constants $h_{eff}/h = n$ assigned to dark matter in TGD framework would allow to circumvent the objection that only objects of length of order Planck length are possible since string tension given by $T = 1/\hbar_{eff}G$ apart from numerical factor could be arbitrary small. This would make possible gravitational bound states as partonic 2-surfaces as structures connected by strings and solve the basic problem of super string theories. This option allows the natural interpretation of M^4 type

vacuum extremals with CP_2 projection, which is Lagrange manifold as good approximations for space-time sheets at macroscopic length scales. String area does not contribute to the Kähler function at all.

Whether also induced spinor fields associated with Kähler-Dirac action and de-localized inside entire space-time surface should be allowed remains an open question: super-conformal symmetry strongly suggests their presence. A possible interpretation for the corresponding spinor modes could be in terms of dark matter, sparticles, and hierarchy of Planck constants.

It is perhaps useful to make clear what TGD is not and also what new TGD can give to physics.

1. TGD is *not* just General Relativity made concrete by using imbeddings: the 4-surface property is absolutely essential for unifying standard model physics with gravitation and to circumvent the incurable conceptual problems of General Relativity. The many-sheeted space-time of TGD gives rise only at macroscopic limit to GRT space-time as a slightly curved Minkowski space. TGD is *not* a Kaluza-Klein theory although color gauge potentials are analogous to gauge potentials in these theories.

TGD space-time is 4-D and its dimension is due to completely unique conformal properties of light-cone boundary and 3-D light-like surfaces implying enormous extension of the ordinary conformal symmetries. Light-like 3-surfaces represent orbits of partonic 2-surfaces and carry fundamental fermions at 1-D boundaries of string world sheets. TGD is *not* obtained by performing Poincare gauging of space-time to introduce gravitation and plagued by profound conceptual problems.

2. TGD is *not* a particular string model although string world sheets emerge in TGD very naturally as loci for spinor modes: their 2-dimensionality makes among other things possible quantum deformation of quantization known to be physically realized in condensed matter, and conjectured in TGD framework to be crucial for understanding the notion of finite measurement resolution. Hierarchy of objects of dimension up to 4 emerge from TGD: this obviously means analogy with branes of super-string models.

TGD is *not* one more item in the collection of string models of quantum gravitation relying on Planck length mystics. Dark matter becomes an essential element of quantum gravitation and quantum coherence in astrophysical scales is predicted just from the assumption that strings connecting partonic 2-surfaces serve are responsible for gravitational bound states.

TGD is *not* a particular string model although AdS/CFT duality of super-string models generalizes due to the huge extension of conformal symmetries and by the identification of WCW gamma matrices as Noether super-charges of super-symplectic algebra having a natural conformal structure.

3. TGD is *not* a gauge theory. In TGD framework the counterparts of also ordinary gauge symmetries are assigned to super-symplectic algebra (and its Yangian), which is a generalization of Kac-Moody algebras rather than gauge algebra and suffers a fractal hierarchy of symmetry breakings defining hierarchy of criticalities. TGD is *not* one more quantum field theory like structure based on path integral formalism: path integral is replaced with functional integral over 3-surfaces, and the notion of classical space-time becomes exact part of the theory. Quantum theory becomes formally a purely classical theory of WCW spinor fields: only state function reduction is something genuinely quantal.
4. TGD view about spinor fields is *not* the standard one. Spinor fields appear at three levels. Spinor modes of the imbedding space are analogs of spinor modes characterizing incoming and outgoing states in quantum field theories. Induced second quantized spinor fields at space-time level are analogs of stringy spinor fields. Their modes are localized by the well-definedness of electro-magnetic charge and by number theoretic arguments at string world sheets. Kähler-Dirac action is fixed by supersymmetry implying that ordinary gamma matrices are replaced by what I call Kähler-Dirac gamma matrices - this something new. WCW spinor fields, which are classical in the sense that they are not second quantized, serve as analogs of fields of string field theory and imply a geometrization of quantum theory.

5. TGD is in some sense an extremely conservative geometrization of entire quantum physics: *no* additional structures such as gauge fields as independent dynamical degrees of freedom are introduced: Kähler geometry and associated spinor structure are enough. “Topological” in TGD should not be understood as an attempt to reduce physics to torsion (see for instance [?]) or something similar. Rather, TGD space-time is topologically non-trivial in all scales and even the visible structures of everyday world represent non-trivial topology of space-time in TGD Universe.
6. Twistor space - or rather, a generalization of twistor approach replacing masslessness in 4-D sense with masslessness in 8-D sense and thus allowing description of also massive particles - emerges as a technical tool, and its Kähler structure is possible only for $H = M^4 \times CP_2$. What is genuinely new is the infinite-dimensional character of the Kähler geometry making it highly unique, and its generalization to p-adic number fields to describe correlates of cognition. Also the hierarchies of Planck constants $h_{eff} = n \times h$ reducing to the quantum criticality of TGD Universe and p-adic length scales and Zero Energy Ontology represent something genuinely new.

The great challenge is to construct a mathematical theory around these physically very attractive ideas and I have devoted the last thirty seven years for the realization of this dream and this has resulted in eight online books about TGD and nine online books about TGD inspired theory of consciousness and of quantum biology.

1.2 Two Vision About TGD And Their Fusion

As already mentioned, TGD can be interpreted both as a modification of general relativity and generalization of string models.

1.2.1 TGD as a Poincare invariant theory of gravitation

The first approach was born as an attempt to construct a Poincare invariant theory of gravitation. Space-time, rather than being an abstract manifold endowed with a pseudo-Riemannian structure, is regarded as a surface in the 8-dimensional space $H = M^4 \times CP_2$, where M^4 denotes Minkowski space and $CP_2 = SU(3)/U(2)$ is the complex projective space of two complex dimensions [A2, A5, A1, A4].

The identification of the space-time as a sub-manifold [A3, A7] of $M^4 \times CP_2$ leads to an exact Poincare invariance and solves the conceptual difficulties related to the definition of the energy-momentum in General Relativity.

It soon however turned out that sub-manifold geometry, being considerably richer in structure than the abstract manifold geometry, leads to a geometrization of all basic interactions. First, the geometrization of the elementary particle quantum numbers is achieved. The geometry of CP_2 explains electro-weak and color quantum numbers. The different H-chiralities of H -spinors correspond to the conserved baryon and lepton numbers. Secondly, the geometrization of the field concept results. The projections of the CP_2 spinor connection, Killing vector fields of CP_2 and of H -metric to four-surface define classical electro-weak, color gauge fields and metric in X^4 .

The choice of H is unique from the condition that TGD has standard model symmetries. Also number theoretical vision selects $H = M^4 \times CP_2$ uniquely. M^4 and CP_2 are also unique spaces allowing twistor space with Kähler structure.

1.2.2 TGD as a generalization of the hadronic string model

The second approach was based on the generalization of the mesonic string model describing mesons as strings with quarks attached to the ends of the string. In the 3-dimensional generalization 3-surfaces correspond to free particles and the boundaries of the 3- surface correspond to partons in the sense that the quantum numbers of the elementary particles reside on the boundaries. Various boundary topologies (number of handles) correspond to various fermion families so that one obtains an explanation for the known elementary particle quantum numbers. This approach leads also to a natural topological description of the particle reactions as topology changes: for instance, two-particle decay corresponds to a decay of a 3-surface to two disjoint 3-surfaces.

This decay vertex does not however correspond to a direct generalization of trouser vertex of string models. Indeed, the important difference between TGD and string models is that the analogs of string world sheet diagrams do not describe particle decays but the propagation of particles via different routes. Particle reactions are described by generalized Feynman diagrams for which 3-D light-like surface describing particle propagating join along their ends at vertices. As 4-manifolds the space-time surfaces are therefore singular like Feynman diagrams as 1-manifolds.

Quite recently, it has turned out that fermionic strings inside space-time surfaces define an exact part of quantum TGD and that this is essential for understanding gravitation in long length scales. Also the analog of AdS/CFT duality emerges in that the Kähler metric can be defined either in terms of Kähler function identifiable as Kähler action assignable to Euclidian space-time regions or Kähler action + string action assignable to Minkowskian regions.

The recent view about construction of scattering amplitudes is very “stringy”. By strong form of holography string world sheets and partonic 2-surfaces provide the data needed to construct scattering amplitudes. Space-time surfaces are however needed to realize quantum-classical correspondence necessary to understand the classical correlates of quantum measurement. There is a huge generalization of the duality symmetry of hadronic string models. Scattering amplitudes can be regarded as sequences of computational operations for the Yangian of super-symplectic algebra. Product and co-product define the basic vertices and realized geometrically as partonic 2-surfaces and algebraically as multiplication for the elements of Yangian identified as super-symplectic Noether charges assignable to strings. Any computational sequences connecting given collections of algebraic objects at the opposite boundaries of causal diamond (CD) produce identical scattering amplitudes.

1.2.3 Fusion of the two approaches via a generalization of the space-time concept

The problem is that the two approaches to TGD seem to be mutually exclusive since the orbit of a particle like 3-surface defines 4-dimensional surface, which differs drastically from the topologically trivial macroscopic space-time of General Relativity. The unification of these approaches forces a considerable generalization of the conventional space-time concept. First, the topologically trivial 3-space of General Relativity is replaced with a “topological condensate” containing matter as particle like 3-surfaces “glued” to the topologically trivial background 3-space by connected sum operation. Secondly, the assumption about connectedness of the 3-space is given up. Besides the “topological condensate” there could be “vapor phase” that is a “gas” of particle like 3-surfaces and string like objects (counterpart of the “baby universes” of GRT) and the non-conservation of energy in GRT corresponds to the transfer of energy between different sheets of the space-time and possibly existence vapour phase.

What one obtains is what I have christened as many-sheeted space-time (see **Fig.** <http://tgdtheory.fi/appfigures/manysheeted.jpg> or **Fig. ??** in the appendix of this book). One particular aspect is topological field quantization meaning that various classical fields assignable to a physical system correspond to space-time sheets representing the classical fields to that particular system. One can speak of the field body of a particular physical system. Field body consists of topological light rays, and electric and magnetic flux quanta. In Maxwell’s theory system does not possess this kind of field identity. The notion of magnetic body is one of the key players in TGD inspired theory of consciousness and quantum biology.

This picture became more detailed with the advent of zero energy ontology (ZEO). The basic notion of ZEO is causal diamond (CD) identified as the Cartesian product of CP_2 and of the intersection of future and past directed light-cones and having scale coming as an integer multiple of CP_2 size is fundamental. CDs form a fractal hierarchy and zero energy states decompose to products of positive and negative energy parts assignable to the opposite boundaries of CD defining the ends of the space-time surface. The counterpart of zero energy state in positive energy ontology is the pair of initial and final states of a physical event, say particle reaction.

At space-time level ZEO means that 3-surfaces are pairs of space-like 3-surfaces at the opposite light-like boundaries of CD. Since the extremals of Kähler action connect these, one can say that by holography the basic dynamical objects are the space-time surface connecting these 3-surfaces. This changes totally the vision about notions like self-organization: self-organization by quantum jumps does not take for a 3-D system but for the entire 4-D field pattern associated with it.

General Coordinate Invariance (GCI) allows to identify the basic dynamical objects as space-like 3-surfaces at the ends of space-time surface at boundaries of CD: this means that space-time surface is analogous to Bohr orbit. An alternative identification is as light-like 3-surfaces at which the signature of the induced metric changes from Minkowskian to Euclidian and interpreted as lines of generalized Feynman diagrams. Also the Euclidian 4-D regions would have similar interpretation. The requirement that the two interpretations are equivalent, leads to a strong form of General Coordinate Invariance. The outcome is effective 2-dimensionality stating that the partonic 2-surfaces identified as intersections of the space-like ends of space-time surface and light-like wormhole throats are the fundamental objects. That only effective 2-dimensionality is in question is due to the effects caused by the failure of strict determinism of Kähler action. In finite length scale resolution these effects can be neglected below UV cutoff and above IR cutoff. One can also speak about strong form of holography.

1.3 Basic Objections

Objections are the most powerful tool in theory building. The strongest objection against TGD is the observation that all classical gauge fields are expressible in terms of four imbedding space coordinates only- essentially CP_2 coordinates. The linear superposition of classical gauge fields taking place independently for all gauge fields is lost. This would be a catastrophe without many-sheeted space-time. Instead of gauge fields, only the effects such as gauge forces are superposed. Particle topologically condenses to several space-time sheets simultaneously and experiences the sum of gauge forces. This transforms the weakness to extreme economy: in a typical unified theory the number of primary field variables is countered in hundreds if not thousands, now it is just four.

Second objection is that TGD space-time is quite too simple as compared to GRT space-time due to the imbeddability to 8-D imbedding space. One can also argue that Poincare invariant theory of gravitation cannot be consistent with General Relativity. The above interpretation allows to understand the relationship to GRT space-time and how Equivalence Principle (EP) follows from Poincare invariance of TGD. The interpretation of GRT space-time is as effective space-time obtained by replacing many-sheeted space-time with Minkowski space with effective metric determined as a sum of Minkowski metric and sum over the deviations of the induced metrics of space-time sheets from Minkowski metric. Poincare invariance suggests strongly classical EP for the GRT limit in long length scales at least. One can consider also other kinds of limits such as the analog of GRT limit for Euclidian space-time regions assignable to elementary particles. In this case deformations of CP_2 metric define a natural starting point and CP_2 indeed defines a gravitational instanton with very large cosmological constant in Einstein-Maxwell theory. Also gauge potentials of standard model correspond classically to superpositions of induced gauge potentials over space-time sheets.

1.3.1 Topological field quantization

Topological field quantization distinguishes between TGD based and more standard - say Maxwellian - notion of field. In Maxwell's fields created by separate systems superpose and one cannot tell which part of field comes from which system except theoretically. In TGD these fields correspond to different space-time sheets and only their effects on test particle superpose. Hence physical systems have well-defined field identifies - field bodies - in particular magnetic bodies.

The notion of magnetic body carrying dark matter with non-standard large value of Planck constant has become central concept in TGD inspired theory of consciousness and living matter, and by starting from various anomalies of biology one ends up to a rather detailed view about the role of magnetic body as intentional agent receiving sensory input from the biological body and controlling it using EEG and its various scaled up variants as a communication tool. Among other things this leads to models for cell membrane, nerve pulse, and EEG.

1.4 P-Adic Variants Of Space-Time Surfaces

There is a further generalization of the space-time concept inspired by p-adic physics forcing a generalization of the number concept through the fusion of real numbers and various p-adic number fields. One might say that TGD space-time is adelic. Also the hierarchy of Planck constants forces

a generalization of the notion of space-time but this generalization can be understood in terms of the failure of strict determinism for Kähler action defining the fundamental variational principle behind the dynamics of space-time surfaces.

A very concise manner to express how TGD differs from Special and General Relativities could be following. Relativity Principle (Poincare Invariance), General Coordinate Invariance, and Equivalence Principle remain true. What is new is the notion of sub-manifold geometry: this allows to realize Poincare Invariance and geometrize gravitation simultaneously. This notion also allows a geometrization of known fundamental interactions and is an essential element of all applications of TGD ranging from Planck length to cosmological scales. Sub-manifold geometry is also crucial in the applications of TGD to biology and consciousness theory.

1.5 The Threads In The Development Of Quantum TGD

The development of TGD has involved several strongly interacting threads: physics as infinite-dimensional geometry; TGD as a generalized number theory, the hierarchy of Planck constants interpreted in terms of dark matter hierarchy, and TGD inspired theory of consciousness. In the following these threads are briefly described.

The theoretical framework involves several threads.

1. Quantum T(opological) G(eometro)D(ynamics) as a classical spinor geometry for infinite-dimensional WCW, p-adic numbers and quantum TGD, and TGD inspired theory of consciousness and of quantum biology have been for last decade of the second millenium the basic three strongly interacting threads in the tapestry of quantum TGD.
2. The discussions with Tony Smith initiated a fourth thread which deserves the name “TGD as a generalized number theory”. The basic observation was that classical number fields might allow a deeper formulation of quantum TGD. The work with Riemann hypothesis made time ripe for realization that the notion of infinite primes could provide, not only a reformulation, but a deep generalization of quantum TGD. This led to a thorough and extremely fruitful revision of the basic views about what the final form and physical content of quantum TGD might be. Together with the vision about the fusion of p-adic and real physics to a larger coherent structure these sub-threads fused to the “physics as generalized number theory” thread.
3. A further thread emerged from the realization that by quantum classical correspondence TGD predicts an infinite hierarchy of macroscopic quantum systems with increasing sizes, that it is not at all clear whether standard quantum mechanics can accommodate this hierarchy, and that a dynamical quantized Planck constant might be necessary and strongly suggested by the failure of strict determinism for the fundamental variational principle. The identification of hierarchy of Planck constants labelling phases of dark matter would be natural. This also led to a solution of a long standing puzzle: what is the proper interpretation of the predicted fractal hierarchy of long ranged classical electro-weak and color gauge fields. Quantum classical correspondences allows only single answer: there is infinite hierarchy of p-adically scaled up variants of standard model physics and for each of them also dark hierarchy. Thus TGD Universe would be fractal in very abstract and deep sense.

The chronology based identification of the threads is quite natural but not logical and it is much more logical to see p-adic physics, the ideas related to classical number fields, and infinite primes as sub-threads of a thread which might be called “physics as a generalized number theory”. In the following I adopt this view. This reduces the number of threads to four.

TGD forces the generalization of physics to a quantum theory of consciousness, and represent TGD as a generalized number theory vision leads naturally to the emergence of p-adic physics as physics of cognitive representations. The eight online books [K20, K12, K8, K27, K16, K26, K25, K15] about TGD and nine online books about TGD inspired theory of consciousness and of quantum biology [K18, K3, K9, K2, K5, K6, K7, K14, K24] are warmly recommended to the interested reader.

1.5.1 Quantum TGD as spinor geometry of World of Classical Worlds

A turning point in the attempts to formulate a mathematical theory was reached after seven years from the birth of TGD. The great insight was “Do not quantize”. The basic ingredients to the new approach have served as the basic philosophy for the attempt to construct Quantum TGD since then and have been the following ones:

1. Quantum theory for extended particles is free(!), classical(!) field theory for a generalized Schrödinger amplitude in the configuration space CH (“world of classical worlds”, WCW) consisting of all possible 3-surfaces in H . “All possible” means that surfaces with arbitrary many disjoint components and with arbitrary internal topology and also singular surfaces topologically intermediate between two different manifold topologies are included. Particle reactions are identified as topology changes [A6, A8, A9]. For instance, the decay of a 3-surface to two 3-surfaces corresponds to the decay $A \rightarrow B + C$. Classically this corresponds to a path of WCW leading from 1-particle sector to 2-particle sector. At quantum level this corresponds to the dispersion of the generalized Schrödinger amplitude localized to 1-particle sector to two-particle sector. All coupling constants should result as predictions of the theory since no nonlinearities are introduced.
2. During years this naive and very rough vision has of course developed a lot and is not anymore quite equivalent with the original insight. In particular, the space-time correlates of Feynman graphs have emerged from theory as Euclidian space-time regions and the strong form of General Coordinate Invariance has led to a rather detailed and in many respects unexpected visions. This picture forces to give up the idea about smooth space-time surfaces and replace space-time surface with a generalization of Feynman diagram in which vertices represent the failure of manifold property. I have also introduced the word “world of classical worlds” (WCW) instead of rather formal “configuration space”. I hope that “WCW” does not induce despair in the reader having tendency to think about the technicalities involved!
3. WCW is endowed with metric and spinor structure so that one can define various metric related differential operators, say Dirac operator, appearing in the field equations of the theory ¹
4. WCW Dirac operator appearing in Super-Virasoro conditions, imbedding space Dirac operator whose modes define the ground states of Super-Virasoro representations, Kähler-Dirac operator at space-time surfaces, and the algebraic variant of M^4 Dirac operator appearing in propagators. The most ambitious dream is that zero energy states correspond to a complete solution basis for the Dirac operator of WCW so that this classical free field theory would dictate M-matrices defined between positive and negative energy parts of zero energy states which form orthonormal rows of what I call U-matrix as a matrix defined between zero energy states. Given M-matrix in turn would decompose to a product of a hermitian square root of density matrix and unitary S-matrix.

M-matrix would define time-like entanglement coefficients between positive and negative energy parts of zero energy states (all net quantum numbers vanish for them) and can be regarded as a hermitian square root of density matrix multiplied by a unitary S-matrix. Quantum theory would be in well-defined sense a square root of thermodynamics. The orthogonality and hermiticity of the M-matrices commuting with S-matrix means that they span infinite-dimensional Lie algebra acting as symmetries of the S-matrix. Therefore quantum TGD would reduce to group theory in well-defined sense.

In fact the Lie algebra of Hermitian M-matrices extends to Kac-Moody type algebra obtained by multiplying hermitian square roots of density matrices with powers of the S-matrix. Also the analog of Yangian algebra involving only non-negative powers of S-matrix is possible and would correspond to a hierarchy of CDs with the temporal distances between tips coming as integer multiples of the CP_2 time.

¹There are four kinds of Dirac operators in TGD. The geometrization of quantum theory requires Kähler metric definable either in terms of Kähler function identified as Kähler action for Euclidian space-time regions or as anti-commutators for WCW gamma matrices identified as conformal Noether super-charges associated with the second quantized modified Dirac action consisting of string world sheet term and possibly also Kähler Dirac action in Minkowskian space-time regions. These two possible definitions reflect a duality analogous to AdS/CFT duality.

The M-matrices associated with CDs are obtained by a discrete scaling from the minimal CD and characterized by integer n are naturally proportional to a representation matrix of scaling: $S(n) = S^n$, where S is unitary S-matrix associated with the minimal CD [K22]. This conforms with the idea about unitary time evolution as exponent of Hamiltonian discretized to integer power of S and represented as scaling with respect to the logarithm of the proper time distance between the tips of CD.

U-matrix elements between M-matrices for various CDs are proportional to the inner products $Tr[S^{-n_1} \circ H^i H^j \circ S^{n_2} \lambda]$, where λ represents unitarily the discrete Lorentz boost relating the moduli of the active boundary of CD and H^i form an orthonormal basis of Hermitian square roots of density matrices. \circ tells that S acts at the active boundary of CD only. It turns out possible to construct a general representation for the U-matrix reducing its construction to that of S-matrix. S-matrix has interpretation as exponential of the Virasoro generator L_{-1} of the Virasoro algebra associated with super-symplectic algebra.

5. By quantum classical correspondence the construction of WCW spinor structure reduces to the second quantization of the induced spinor fields at space-time surface. The basic action is so called modified Dirac action (or Kähler-Dirac action) in which gamma matrices are replaced with the modified (Kähler-Dirac) gamma matrices defined as contractions of the canonical momentum currents with the imbedding space gamma matrices. In this manner one achieves super-conformal symmetry and conservation of fermionic currents among other things and consistent Dirac equation. The Kähler-Dirac gamma matrices define as anti-commutators effective metric, which might provide geometrization for some basic observables of condensed matter physics. One might also talk about bosonic emergence in accordance with the prediction that the gauge bosons and graviton are expressible in terms of bound states of fermion and anti-fermion.
6. An important result relates to the notion of induced spinor connection. If one requires that spinor modes have well-defined em charge, one must assume that the modes in the generic situation are localized at 2-D surfaces - string world sheets or perhaps also partonic 2-surfaces - at which classical W boson fields vanish. Covariantly constant right handed neutrino generating super-symmetries forms an exception. The vanishing of also Z^0 field is possible for Kähler-Dirac action and should hold true at least above weak length scales. This implies that string model in 4-D space-time becomes part of TGD. Without these conditions classical weak fields can vanish above weak scale only for the GRT limit of TGD for which gauge potentials are sums over those for space-time sheets.

The localization simplifies enormously the mathematics and one can solve exactly the Kähler-Dirac equation for the modes of the induced spinor field just like in super string models.

At the light-like 3-surfaces at which the signature of the induced metric changes from Euclidian to Minkowskian so that $\sqrt{g_4}$ vanishes one can pose the condition that the algebraic analog of massless Dirac equation is satisfied by the nodes so that Kähler-Dirac action gives massless Dirac propagator localizable at the boundaries of the string world sheets.

The evolution of these basic ideas has been rather slow but has gradually led to a rather beautiful vision. One of the key problems has been the definition of Kähler function. Kähler function is Kähler action for a preferred extremal assignable to a given 3-surface but what this preferred extremal is? The obvious first guess was as absolute minimum of Kähler action but could not be proven to be right or wrong. One big step in the progress was boosted by the idea that TGD should reduce to almost topological QFT in which braids would replace 3-surfaces in finite measurement resolution, which could be inherent property of the theory itself and imply discretization at partonic 2-surfaces with discrete points carrying fermion number.

It took long time to realize that there is no discretization in 4-D sense - this would lead to difficulties with basic symmetries. Rather, the discretization occurs for the parameters characterizing co-dimension 2 objects representing the information about space-time surface so that they belong to some algebraic extension of rationals. These 2-surfaces - string world sheets and partonic 2-surfaces - are genuine physical objects rather than a computational approximation. Physics itself approximates itself, one might say! This is of course nothing but strong form of holography.

1. TGD as almost topological QFT vision suggests that Kähler action for preferred extremals reduces to Chern-Simons term assigned with space-like 3-surfaces at the ends of space-time (recall the notion of causal diamond (CD)) and with the light-like 3-surfaces at which the signature of the induced metric changes from Minkowskian to Euclidian. Minkowskian and Euclidian regions would give at wormhole throats the same contribution apart from coefficients and in Minkowskian regions the $\sqrt{g_4}$ factor coming from metric would be imaginary so that one would obtain sum of real term identifiable as Kähler function and imaginary term identifiable as the ordinary Minkowskian action giving rise to interference effects and stationary phase approximation central in both classical and quantum field theory.

Imaginary contribution - the presence of which I realized only after 33 years of TGD - could also have topological interpretation as a Morse function. On physical side the emergence of Euclidian space-time regions is something completely new and leads to a dramatic modification of the ideas about black hole interior.

2. The manner to achieve the reduction to Chern-Simons terms is simple. The vanishing of Coulomb contribution to Kähler action is required and is true for all known extremals if one makes a general ansatz about the form of classical conserved currents. The so called weak form of electric-magnetic duality defines a boundary condition reducing the resulting 3-D terms to Chern-Simons terms. In this manner almost topological QFT results. But only “almost” since the Lagrange multiplier term forcing electric-magnetic duality implies that Chern-Simons action for preferred extremals depends on metric.

1.5.2 TGD as a generalized number theory

Quantum T(opological)D(ynamics) as a classical spinor geometry for infinite-dimensional configuration space (“world of classical worlds”, WCW), p-adic numbers and quantum TGD, and TGD inspired theory of consciousness, have been for last ten years the basic three strongly interacting threads in the tapestry of quantum TGD. The fourth thread deserves the name “TGD as a generalized number theory”. It involves three separate threads: the fusion of real and various p-adic physics to a single coherent whole by requiring number theoretic universality discussed already, the formulation of quantum TGD in terms of hyper-counterparts of classical number fields identified as sub-spaces of complexified classical number fields with Minkowskian signature of the metric defined by the complexified inner product, and the notion of infinite prime.

1. *p-Adic TGD and fusion of real and p-adic physics to single coherent whole*

The p-adic thread emerged for roughly ten years ago as a dim hunch that p-adic numbers might be important for TGD. Experimentation with p-adic numbers led to the notion of canonical identification mapping reals to p-adics and vice versa. The breakthrough came with the successful p-adic mass calculations using p-adic thermodynamics for Super-Virasoro representations with the super-Kac-Moody algebra associated with a Lie-group containing standard model gauge group. Although the details of the calculations have varied from year to year, it was clear that p-adic physics reduces not only the ratio of proton and Planck mass, the great mystery number of physics, but all elementary particle mass scales, to number theory if one assumes that primes near prime powers of two are in a physically favored position. Why this is the case, became one of the key puzzles and led to a number of arguments with a common gist: evolution is present already at the elementary particle level and the primes allowed by the p-adic length scale hypothesis are the fittest ones.

It became very soon clear that p-adic topology is not something emerging in Planck length scale as often believed, but that there is an infinite hierarchy of p-adic physics characterized by p-adic length scales varying to even cosmological length scales. The idea about the connection of p-adics with cognition motivated already the first attempts to understand the role of the p-adics and inspired “Universe as Computer” vision but time was not ripe to develop this idea to anything concrete (p-adic numbers are however in a central role in TGD inspired theory of consciousness). It became however obvious that the p-adic length scale hierarchy somehow corresponds to a hierarchy of intelligences and that p-adic prime serves as a kind of intelligence quotient. Ironically, the almost obvious idea about p-adic regions as cognitive regions of space-time providing cognitive representations for real regions had to wait for almost a decade for the access into my consciousness.

In string model context one tries to reduce the physics to Planck scale. The price is the inability to say anything about physics in long length scales. In TGD p-adic physics takes care of this shortcoming by predicting the physics also in long length scales.

There were many interpretational and technical questions crying for a definite answer.

1. What is the relationship of p-adic non-determinism to the classical non-determinism of the basic field equations of TGD? Are the p-adic space-time region genuinely p-adic or does p-adic topology only serve as an effective topology? If p-adic physics is direct image of real physics, how the mapping relating them is constructed so that it respects various symmetries? Is the basic physics p-adic or real (also real TGD seems to be free of divergences) or both? If it is both, how should one glue the physics in different number field together to get *the* Physics? Should one perform p-adicization also at the level of the WCW? Certainly the p-adicization at the level of super-conformal representation is necessary for the p-adic mass calculations.
2. Perhaps the most basic and most irritating technical problem was how to precisely define p-adic definite integral which is a crucial element of any variational principle based formulation of the field equations. Here the frustration was not due to the lack of solution but due to the too large number of solutions to the problem, a clear symptom for the sad fact that clever inventions rather than real discoveries might be in question. Quite recently I however learned that the problem of making sense about p-adic integration has been for decades central problem in the frontier of mathematics and a lot of profound work has been done along same intuitive lines as I have proceeded in TGD framework. The basic idea is certainly the notion of algebraic continuation from the world of rationals belonging to the intersection of real world and various p-adic worlds.

Despite various uncertainties, the number of the applications of the poorly defined p-adic physics has grown steadily and the applications turned out to be relatively stable so that it was clear that the solution to these problems must exist. It became only gradually clear that the solution of the problems might require going down to a deeper level than that represented by reals and p-adics.

The key challenge is to fuse various p-adic physics and real physics to single larger structures. This has inspired a proposal for a generalization of the notion of number field by fusing real numbers and various p-adic number fields and their extensions along rationals and possible common algebraic numbers. This leads to a generalization of the notions of imbedding space and space-time concept and one can speak about real and p-adic space-time sheets. One can talk about adelic space-time, imbedding space, and WCW.

The notion of p-adic manifold [K28] identified as p-adic space-time surface solving p-adic analogs of field equations and having real space-time sheet as chart map provided a possible solution of the basic challenge of relating real and p-adic classical physics. One can also speak of real space-time surfaces having p-adic space-time surfaces as chart maps (cognitive maps, “thought bubbles”). Discretization required having interpretation in terms of finite measurement resolution is unavoidable in this approach and this leads to problems with symmetries: canonical identification does not commute with symmetries.

It is now clear that much more elegant approach based on abstraction exists [K29]. The map of real preferred extremals to p-adic ones is not induced from a local correspondence between points but is global. Discretization occurs only for the parameters characterizing string world sheets and partonic 2-surfaces so that they belong to some algebraic extension of rationals. Restriction to these 2-surfaces is possible by strong form of holography. Adelization providing number theoretical universality reduces to algebraic continuation for the amplitudes from this intersection of reality and various p-adicities - analogous to a back of a book - to various number fields. There are no problems with symmetries but canonical identification is needed: various group invariant of the amplitude are mapped by canonical identification to various p-adic number fields. This is nothing but a generalization of the mapping of the p-adic mass squared to its real counterpart in p-adic mass calculations.

This leads to surprisingly detailed predictions and far reaching conjectures. For instance, the number theoretic generalization of entropy concept allows negentropic entanglement central for the applications to living matter (see **Fig.** <http://tgdtheory.fi/appfigures/cat.jpg> or **Fig. ??** in the appendix of this book). One can also understand how preferred p-adic primes emerge as so called ramified primes of algebraic extension of rationals in question and characterizing string world

sheets and partonic 2-surfaces. Also a generalization of p-adic length scale hypothesis emerges from NMP.

The characteristic non-determinism of the p-adic differential equations suggests strongly that p-adic regions correspond to “mind stuff”, the regions of space-time where cognitive representations reside. This interpretation implies that p-adic physics is physics of cognition. Since Nature is probably a brilliant simulator of Nature, the natural idea is to study the p-adic physics of the cognitive representations to derive information about the real physics. This view encouraged by TGD inspired theory of consciousness clarifies difficult interpretational issues and provides a clear interpretation for the predictions of p-adic physics.

2. The role of classical number fields

The vision about the physical role of the classical number fields relies on certain speculative questions inspired by the idea that space-time dynamics could be reduced to associativity or co-associativity condition. Associativity means here associativity of tangent spaces of space-time region and co-associativity associativity of normal spaces of space-time region.

1. Could space-time surfaces X^4 be regarded as associative or co-associative (“quaternionic” is equivalent with “associative”) surfaces of H endowed with octonionic structure in the sense that tangent space of space-time surface would be associative (co-associative with normal space associative) sub-space of octonions at each point of X^4 [K17]. This is certainly possible and an interesting conjecture is that the preferred extremals of Kähler action include associative and co-associative space-time regions.
2. Could the notion of compactification generalize to that of number theoretic compactification in the sense that one can map associative (co-associative) surfaces of M^8 regarded as octonionic linear space to surfaces in $M^4 \times CP_2$ [K17] ? This conjecture - $M^8 - H$ duality - would give for $M^4 \times CP_2$ deep number theoretic meaning. CP_2 would parametrize associative planes of octonion space containing fixed complex plane $M^2 \subset M^8$ and CP_2 point would thus characterize the tangent space of $X^4 \subset M^8$. The point of M^4 would be obtained by projecting the point of $X^4 \subset M^8$ to a point of M^4 identified as tangent space of X^4 . This would guarantee that the dimension of space-time surface in H would be four. The conjecture is that the preferred extremals of Kähler action include these surfaces.
3. $M^8 - H$ duality can be generalized to a duality $H \rightarrow H$ if the images of the associative surface in M^8 is associative surface in H . One can start from associative surface of H and assume that it contains the preferred M^2 tangent plane in 8-D tangent space of H or integrable distribution $M^2(x)$ of them, and its points to H by mapping M^4 projection of H point to itself and associative tangent space to CP_2 point. This point need not be the original one! If the resulting surface is also associative, one can iterate the process indefinitely. WCW would be a category with one object.
4. G_2 defines the automorphism group of octonions, and one might hope that the maps of octonions to octonions such that the action of Jacobian in the tangent space of associative or co-associative surface reduces to that of G_2 could produce new associative/co-associative surfaces. The action of G_2 would be analogous to that of gauge group.
5. One can also ask whether the notions of commutativity and co-commutativity could have physical meaning. The well-definedness of em charge as quantum number for the modes of the induced spinor field requires their localization to 2-D surfaces (right-handed neutrino is an exception) - string world sheets and partonic 2-surfaces. This can be possible only for Kähler action and could have commutativity and co-commutativity as a number theoretic counterpart. The basic vision would be that the dynamics of Kähler action realizes number theoretical geometrical notions like associativity and commutativity and their co-notions.

The notion of number theoretic compactification stating that space-time surfaces can be regarded as surfaces of either M^8 or $M^4 \times CP_2$. As surfaces of M^8 identifiable as space of hyper-octonions they are hyper-quaternionic or co-hyper-quaternionic- and thus maximally associative or co-associative. This means that their tangent space is either hyper-quaternionic plane of M^8 or an orthogonal complement of such a plane. These surface can be mapped in natural manner to

surfaces in $M^4 \times CP_2$ [K17] provided one can assign to each point of tangent space a hyper-complex plane $M^2(x) \subset M^4 \subset M^8$. One can also speak about $M^8 - H$ duality.

This vision has very strong predictive power. It predicts that the preferred extremals of Kähler action correspond to either hyper-quaternionic or co-hyper-quaternionic surfaces such that one can assign to tangent space at each point of space-time surface a hyper-complex plane $M^2(x) \subset M^4$. As a consequence, the M^4 projection of space-time surface at each point contains $M^2(x)$ and its orthogonal complement. These distributions are integrable implying that space-time surface allows dual slicings defined by string world sheets Y^2 and partonic 2-surfaces X^2 . The existence of this kind of slicing was earlier deduced from the study of extremals of Kähler action and christened as Hamilton-Jacobi structure. The physical interpretation of $M^2(x)$ is as the space of non-physical polarizations and the plane of local 4-momentum.

Number theoretical compactification has inspired large number of conjectures. This includes dual formulations of TGD as Minkowskian and Euclidian string model type theories, the precise identification of preferred extremals of Kähler action as extremals for which second variation vanishes (at least for deformations representing dynamical symmetries) and thus providing space-time correlate for quantum criticality, the notion of number theoretic braid implied by the basic dynamics of Kähler action and crucial for precise construction of quantum TGD as almost-topological QFT, the construction of WCW metric and spinor structure in terms of second quantized induced spinor fields with modified Dirac action defined by Kähler action realizing the notion of finite measurement resolution and a connection with inclusions of hyper-finite factors of type II_1 about which Clifford algebra of WCW represents an example.

The two most important number theoretic conjectures relate to the preferred extremals of Kähler action. The general idea is that classical dynamics for the preferred extremals of Kähler action should reduce to number theory: space-time surfaces should be either associative or co-associative in some sense.

Associativity (co-associativity) would be that tangent (normal) spaces of space-time surfaces associative (co-associative) in some sense and thus quaternionic (co-quaternionic). This can be formulated in two manners.

1. One can introduce octonionic tangent space basis by assigning to the “free” gamma matrices octonion basis or in terms of octonionic representation of the imbedding space gamma matrices possible in dimension $D = 8$.
2. Associativity (quaternionicity) would state that the projections of octonionic basic vectors or induced gamma matrices basis to the space-time surface generates associative (quaternionic) sub-algebra at each space-time point. Co-associativity is defined in analogous manner and can be expressed in terms of the components of second fundamental form.
3. For gamma matrix option induced rather than Kähler-Dirac gamma matrices must be in question since Kähler-Dirac gamma matrices can span lower than 4-dimensional space and are not parallel to the space-time surfaces as imbedding space vectors.

3. Infinite primes

The discovery of the hierarchy of infinite primes and their correspondence with a hierarchy defined by a repeatedly second quantized arithmetic quantum field theory gave a further boost for the speculations about TGD as a generalized number theory.

After the realization that infinite primes can be mapped to polynomials possibly representable as surfaces geometrically, it was clear how TGD might be formulated as a generalized number theory with infinite primes forming the bridge between classical and quantum such that real numbers, p-adic numbers, and various generalizations of p-adics emerge dynamically from algebraic physics as various completions of the algebraic extensions of rational (hyper-)quaternions and (hyper-)octonions. Complete algebraic, topological and dimensional democracy would characterize the theory.

The infinite primes at the first level of hierarchy, which represent analogs of bound states, can be mapped to irreducible polynomials, which in turn characterize the algebraic extensions of rationals defining a hierarchy of algebraic physics continuable to real and p-adic number fields. The products of infinite primes in turn define more general algebraic extensions of rationals. The

interesting question concerns the physical interpretation of the higher levels in the hierarchy of infinite primes and integers mappable to polynomials of $n > 1$ variables.

1.6 Hierarchy Of Planck Constants And Dark Matter Hierarchy

By quantum classical correspondence space-time sheets can be identified as quantum coherence regions. Hence the fact that they have all possible size scales more or less unavoidably implies that Planck constant must be quantized and have arbitrarily large values. If one accepts this then also the idea about dark matter as a macroscopic quantum phase characterized by an arbitrarily large value of Planck constant emerges naturally as does also the interpretation for the long ranged classical electro-weak and color fields predicted by TGD. Rather seldom the evolution of ideas follows simple linear logic, and this was the case also now. In any case, this vision represents the fifth, relatively new thread in the evolution of TGD and the ideas involved are still evolving.

1.6.1 Dark matter as large \hbar phases

D. Da Rocha and Laurent Nottale [E1] have proposed that Schrödinger equation with Planck constant \hbar replaced with what might be called gravitational Planck constant $\hbar_{gr} = \frac{GmM}{v_0}$ ($\hbar = c = 1$). v_0 is a velocity parameter having the value $v_0 = 144.7 \pm .7$ km/s giving $v_0/c = 4.6 \times 10^{-4}$. This is rather near to the peak orbital velocity of stars in galactic halos. Also subharmonics and harmonics of v_0 seem to appear. The support for the hypothesis coming from empirical data is impressive.

Nottale and Da Rocha believe that their Schrödinger equation results from a fractal hydrodynamics. Many-sheeted space-time however suggests that astrophysical systems are at some levels of the hierarchy of space-time sheets macroscopic quantum systems. The space-time sheets in question would carry dark matter.

Nottale's hypothesis would predict a gigantic value of \hbar_{gr} . Equivalence Principle and the independence of gravitational Compton length on mass m implies however that one can restrict the values of mass m to masses of microscopic objects so that \hbar_{gr} would be much smaller. Large \hbar_{gr} could provide a solution of the black hole collapse (IR catastrophe) problem encountered at the classical level. The resolution of the problem inspired by TGD inspired theory of living matter is that it is the dark matter at larger space-time sheets which is quantum coherent in the required time scale [K13].

It is natural to assign the values of Planck constants postulated by Nottale to the space-time sheets mediating gravitational interaction and identifiable as magnetic flux tubes (quanta) possibly carrying monopole flux and identifiable as remnants of cosmic string phase of primordial cosmology. The magnetic energy of these flux quanta would correspond to dark energy and magnetic tension would give rise to negative "pressure" forcing accelerate cosmological expansion. This leads to a rather detailed vision about the evolution of stars and galaxies identified as bubbles of ordinary and dark matter inside magnetic flux tubes identifiable as dark energy.

Certain experimental findings suggest the identification $\hbar_{eff} = n \times \hbar_{gr}$. The large value of \hbar_{gr} can be seen as a manner to reduce the string tension of fermionic strings so that gravitational (in fact all!) bound states can be described in terms of strings connecting the partonic 2-surfaces defining particles (analogous to AdS/CFT description). The values $\hbar_{eff}/\hbar = n$ can be interpreted in terms of a hierarchy of breakings of super-conformal symmetry in which the super-conformal generators act as gauge symmetries only for a sub-algebras with conformal weights coming as multiples of n . Macroscopic quantum coherence in astrophysical scales is implied. If also Kähler-Dirac action is present, part of the interior degrees of freedom associated with the Kähler-Dirac part of conformal algebra become physical. A possible is that fermionic oscillator operators generate super-symmetries and sparticles correspond almost by definition to dark matter with $\hbar_{eff}/\hbar = n > 1$. One implication would be that at least part if not all gravitons would be dark and be observed only through their decays to ordinary high frequency graviton ($E = \hbar f_{high} = \hbar_{eff} f_{low}$) of bunch of n low energy gravitons.

1.6.2 *Hierarchy of Planck constants from the anomalies of neuroscience and biology*

The quantal ELF effects of ELF em fields on vertebrate brain have been known since seventies. ELF em fields at frequencies identifiable as cyclotron frequencies in magnetic field whose intensity is about 2/5 times that of Earth for biologically important ions have physiological effects and affect also behavior. What is intriguing that the effects are found only in vertebrates (to my best knowledge). The energies for the photons of ELF em fields are extremely low - about 10^{-10} times lower than thermal energy at physiological temperatures- so that quantal effects are impossible in the framework of standard quantum theory. The values of Planck constant would be in these situations large but not gigantic.

This inspired the hypothesis that these photons correspond to so large a value of Planck constant that the energy of photons is above the thermal energy. The proposed interpretation was as dark photons and the general hypothesis was that dark matter corresponds to ordinary matter with non-standard value of Planck constant. If only particles with the same value of Planck constant can appear in the same vertex of Feynman diagram, the phases with different value of Planck constant are dark relative to each other. The phase transitions changing Planck constant can however make possible interactions between phases with different Planck constant but these interactions do not manifest themselves in particle physics. Also the interactions mediated by classical fields should be possible. Dark matter would not be so dark as we have used to believe.

The hypothesis $h_{eff} = h_{gr}$ - at least for microscopic particles - implies that cyclotron energies of charged particles do not depend on the mass of the particle and their spectrum is thus universal although corresponding frequencies depend on mass. In bio-applications this spectrum would correspond to the energy spectrum of bio-photons assumed to result from dark photons by h_{eff} reducing phase transition and the energies of bio-photons would be in visible and UV range associated with the excitations of bio-molecules.

Also the anomalies of biology (see for instance [K10, K11, K23]) support the view that dark matter might be a key player in living matter.

1.6.3 *Does the hierarchy of Planck constants reduce to the vacuum degeneracy of Kähler action?*

This starting point led gradually to the recent picture in which the hierarchy of Planck constants is postulated to come as integer multiples of the standard value of Planck constant. Given integer multiple $\hbar = n\hbar_0$ of the ordinary Planck constant \hbar_0 is assigned with a multiple singular covering of the imbedding space [K4]. One ends up to an identification of dark matter as phases with non-standard value of Planck constant having geometric interpretation in terms of these coverings providing generalized imbedding space with a book like structure with pages labelled by Planck constants or integers characterizing Planck constant. The phase transitions changing the value of Planck constant would correspond to leakage between different sectors of the extended imbedding space. The question is whether these coverings must be postulated separately or whether they are only a convenient auxiliary tool.

The simplest option is that the hierarchy of coverings of imbedding space is only effective. Many-sheeted coverings of the imbedding space indeed emerge naturally in TGD framework. The huge vacuum degeneracy of Kähler action implies that the relationship between gradients of the imbedding space coordinates and canonical momentum currents is many-to-one: this was the very fact forcing to give up all the standard quantization recipes and leading to the idea about physics as geometry of the “world of classical worlds”. If one allows space-time surfaces for which all sheets corresponding to the same values of the canonical momentum currents are present, one obtains effectively many-sheeted covering of the imbedding space and the contributions from sheets to the Kähler action are identical. If all sheets are treated effectively as one and the same sheet, the value of Planck constant is an integer multiple of the ordinary one. A natural boundary condition would be that at the ends of space-time at future and past boundaries of causal diamond containing the space-time surface, various branches co-incide. This would raise the ends of space-time surface in special physical role.

A more precise formulation is in terms of presence of large number of space-time sheets connecting given space-like 3-surfaces at the opposite boundaries of causal diamond. Quantum criticality presence of vanishing second variations of Kähler action and identified in terms of conformal invari-

ance broken down to to sub-algebras of super-conformal algebras with conformal weights divisible by integer n is highly suggestive notion and would imply that n sheets of the effective covering are actually conformal equivalence classes of space-time sheets with same Kähler action and same values of conserved classical charges (see **Fig.** <http://tgdtheory.fi/appfigures/planckhierarchy.jpg> or **Fig. ??** the appendix of this book). n would naturally correspond the value of h_{eff} and its factors negentropic entanglement with unit density matrix would be between the n sheets of two coverings of this kind. p-Adic prime would be largest prime power factor of n .

1.6.4 Dark matter as a source of long ranged weak and color fields

Long ranged classical electro-weak and color gauge fields are unavoidable in TGD framework. The smallness of the parity breaking effects in hadronic, nuclear, and atomic length scales does not however seem to allow long ranged electro-weak gauge fields. The problem disappears if long range classical electro-weak gauge fields are identified as space-time correlates for massless gauge fields created by dark matter. Also scaled up variants of ordinary electro-weak particle spectra are possible. The identification explains chiral selection in living matter and unbroken $U(2)_{ew}$ invariance and free color in bio length scales become characteristics of living matter and of bio-chemistry and bio-nuclear physics.

The recent view about the solutions of Kähler- Dirac action assumes that the modes have a well-defined em charge and this implies that localization of the modes to 2-D surfaces (right-handed neutrino is an exception). Classical W boson fields vanish at these surfaces and also classical Z^0 field can vanish. The latter would guarantee the absence of large parity breaking effects above intermediate boson scale scaling like h_{eff} .

1.7 Twistors And TGD

8-dimensional generalization of ordinary twistors is highly attractive approach to TGD [K19]. The reason is that M^4 and CP_2 are completely exceptional in the sense that they are the only 4-D manifolds allowing twistor space with Kähler structure. The twistor space of $M^4 \times CP_2$ is Cartesian product of those of M^4 and CP_2 . The obvious idea is that space-time surfaces allowing twistor structure if they are orientable are representable as surfaces in H such that the properly induced twistor structure co-incides with the twistor structure defined by the induced metric. This condition would define the dynamics, and the conjecture is that this dynamics is equivalent with the identification of space-time surfaces as preferred extremals of Kähler action. The dynamics of space-time surfaces would be lifted to the dynamics of twistor spaces, which are sphere bundles over space-time surfaces. What is remarkable that the powerful machinery of complex analysis becomes available.

The condition that the basic formulas for the twistors in M^8 serving as tangent space of imbedding space generalize. This is the case if one introduces octonionic sigma matrices allowing twistor representation of 8-momentum serving as dual for four-momentum and color quantum numbers. The conditions that octonionic spinors are equivalent with ordinary requires that the induced gamma matrices generate quaternionic sub-algebra at given point of string world sheet. This is however not enough: the charge matrices defined by sigma matrices can also break associativity and induced gauge fields must vanish: the CP_2 projection of string world sheet would be one-dimensional at most. This condition is symplectically invariant. Note however that for the interior dynamics of induced spinor fields octonionic representations of Clifford algebra cannot be equivalent with the ordinary one.

One can assign 4-momentum both to the spinor harmonics of the imbedding space representing ground states of superconformal representations and to light-like boundaries of string world sheets at the orbits of partonic 2-surfaces. The two four-momenta should be identical by quantum classical correspondence: this is nothing but a concretization of Equivalence Principle. Also a connection with string model emerges.

Twistor approach developed rapidly during years. Witten's twistor string theory generalizes: the most natural counterpart of Witten's twistor strings is partonic 2-surface. The notion of positive Grassmannian has emerged and TGD provides a possible generalization and number theoretic interpretation of this notion. TGD generalizes the observation that scattering amplitudes in twistor Grassmann approach correspond to representations for permutations. Since 2-vertex is the

only fermionic vertex in TGD, OZI rules for fermions generalizes, and scattering amplitudes are representations for braidings. Braid interpretation gives further support for the conjecture that non-planar diagrams can be reduced to ordinary ones by a procedure analogous to the construction of braid (knot) invariants by gradual un-braiding (un-knotting).

2 Bird's Eye Of View About The Topics Of The Book

The topics of this book are the purely geometric aspects of the vision about physics as an infinite-dimensional Kähler geometry of the “world of classical worlds”, with classical world identified either as light-like 3-D surface of the unique Bohr orbit like 4-surface traversing through it. The non-determinism of Kähler action forces to generalize the notion of 3-surface so that unions of space-like surfaces with time like separations must be allowed. Zero energy ontology allows to formulate this picture elegantly in terms of causal diamonds defined as intersections of future and past directed light-cones. Also a geometric realization of coupling constant evolution and finite measurement resolution emerges.

There are two separate tasks involved.

1. Provide WCW of 3-surfaces with Kähler geometry which is consistent with 4-dimensional general coordinate invariance so that the metric is Diff^4 degenerate. General coordinate invariance implies that the definition of metric must assign to a given light-like 3-surface X^3 a 4-surface as a kind of Bohr orbit $X^4(X^3)$.
2. Provide the WCW with a spinor structure. The great idea is to identify WCW gamma matrices in terms of super algebra generators expressible using second quantized fermionic oscillator operators for induced free spinor fields at the space-time surface assignable to a given 3-surface. The isometry generators and contractions of Killing vectors with gamma matrices would thus form a generalization of Super Kac-Moody algebra.

The condition of mathematical existence poses surprisingly strong conditions on WCW metric and spinor structure.

1. From the experience with loop spaces one can expect that there is no hope about existence of well-defined Riemann connection unless this space is union of infinite-dimensional symmetric spaces with constant curvature metric and simple considerations requires that vacuum Einstein equations are satisfied by each component in the union. The coordinates labeling these symmetric spaces are zero modes having interpretation as genuinely classical variables which do not quantum fluctuate since they do not contribute to the line element of the WCW.
2. The construction of the Kähler structure involves also the identification of complex structure. Direct construction of Kähler function as action associated with a preferred Bohr orbit like extremal for some physically motivated action leads to a unique result. Second approach is group theoretical and is based on a direct guess of isometries of the infinite-dimensional symmetric space formed by 3-surfaces with fixed values of zero modes. The group of isometries is generalization of Kac-Moody group obtained by replacing finite-dimensional Lie group with the group of symplectic transformations of $\delta M_+^4 \times CP_2$, where δM_+^4 is the boundary of 4-dimensional future light-cone. A crucial role is played by the generalized conformal invariance assignable to light-like 3-surfaces and to the boundaries of causal diamond. Contrary to the original belief, the coset construction does not provide a realization of Equivalence Principle at quantum level. The proper realization of EP at quantum level seems to be based on the identification of classical Noether charges in Cartan algebra with the eigenvalues of their quantum counterparts assignable to Kähler-Dirac action. At classical level EP follows at GRT limit obtained by lumping many-sheeted space-time to M^4 with effective metric satisfying Einstein's equations as a reflection of the underlying Poincare invariance.
3. Fermionic statistics and quantization of spinor fields can be realized in terms of WCW spinors structure. Quantum criticality and the idea about space-time surfaces as analogs of Bohr orbits have served as basic guiding lines of Quantum TGD. These notions can be formulated more precisely in terms of the modified Dirac equation for induced spinor fields allowing also

realization of super-conformal symmetries and quantum gravitational holography. A rather detailed view about how WCW Kähler function emerges as Dirac determinant allowing a tentative identification of the preferred extremals of Kähler action as surface for which second variation of Kähler action vanishes for some deformations of the surface. The catastrophe theoretic analog for quantum critical space-time surfaces are the points of space spanned by behavior and control variables at which the determinant defined by the second derivatives of potential function with respect to behavior variables vanishes. Number theoretic vision leads to rather detailed view about preferred extremals of Kähler action. In particular, preferred extremals should be what I have dubbed as hyper-quaternionic surfaces. It is still an open question whether this characterization is equivalent with quantum criticality or not.

3 Sources

The eight online books about TGD [K20, K12, K27, K16, K8, K26, K25, K15] and nine online books about TGD inspired theory of consciousness and quantum biology [K18, K3, K9, K2, K5, K6, K7, K14, K24] are warmly recommended for the reader willing to get overall view about what is involved.

My homepage (<http://www.tgdtheory.com/curri.html>) contains a lot of material about TGD. In particular, there is summary about TGD and its applications using CMAP representation serving also as a TGD glossary [L1, L2] (see <http://tgdtheory.fi/cmaphtml.html> and <http://tgdtheory.fi/tgdglossary.pdf>).

I have published articles about TGD and its applications to consciousness and living matter in *Journal of Non-Locality* (<http://journals.sfu.ca/jnonlocality/index.php/jnonlocality>) founded by Lian Sidorov and in *Prespacetime Journal* (<http://prespacetime.com>), *Journal of Consciousness Research and Exploration* (<https://www.createspace.com/4185546>), and *DNA Decipher Journal* (<http://dnadecipher.com>), all of them founded by Huping Hu. One can find the list about the articles published at <http://www.tgdtheory.com/curri.html>. I am grateful for these far-sighted people for providing a communication channel, whose importance one cannot overestimate.

4 The contents of the book

In the following abstracts of various chapters of the book are given in order to provide overall view.

4.1 Identification of the Configuration Space Kähler Function

There are two basic approaches to quantum TGD. The first approach, which is discussed in this chapter, is a generalization of Einstein's geometrization program of physics to an infinite-dimensional context. Second approach is based on the identification of physics as a generalized number theory. The first approach relies on the vision of quantum physics as infinite-dimensional Kähler geometry for the "world of classical worlds" (WCW) identified as the space of 3-surfaces in in certain 8-dimensional space.

There are three separate manners to meet the challenge of constructing WCW Kähler geometry and spinor structure. The first approach relies on direct guess of Kähler function. Second approach relies on the construction of Kähler form and metric utilizing the huge symmetries of the geometry needed to guarantee the mathematical existence of Riemann connection. The third approach relies on the construction of spinor structure based on the hypothesis that complexified WCW gamma matrices are representable as linear combinations of fermionic oscillator operator for second quantized free spinor fields at space-time surface and on the geometrization of super-conformal symmetries in terms of WCW spinor structure.

In this chapter the proposal for Kähler function based on the requirement of 4-dimensional General Coordinate Invariance implying that its definition must assign to a given 3-surface a unique space-time surface. Quantum classical correspondence requires that this surface is a preferred extremal of some some general coordinate invariant action, and so called Kähler action is a unique candidate in this respect. The preferred extremal has in positive energy ontology interpretation as an analog of Bohr orbit so that classical physics becomes and exact part of WCW geometry

and therefore also quantum physics. In zero energy ontology (ZEO) it is not clear whether this interpretation can be preserved except for maxima of Kähler function.

The basic challenge is the explicit identification of WCW Kähler function K . Two assumptions lead to the identification of K as a sum of Chern-Simons type terms associated with the ends of causal diamond and with the light-like wormhole throats at which the signature of the induced metric changes. The first assumption is the weak form of electric magnetic duality. Second assumption is that the Kähler current for preferred extremals satisfies the condition $j_K \wedge dj_K = 0$ implying that the flow parameter of the flow lines of j_K defines a global space-time coordinate. This would mean that the vision about reduction to almost topological QFT would be realized.

Second challenge is the understanding of the space-time correlates of quantum criticality. Electric-magnetic duality helps considerably here. The realization that the hierarchy of Planck constant realized in terms of coverings of the imbedding space follows from basic quantum TGD leads to a further understanding. The extreme non-linearity of canonical momentum densities as functions of time derivatives of the imbedding space coordinates implies that the correspondence between these two variables is not 1-1 so that it is natural to introduce coverings of $CD \times CP_2$. This leads also to a precise geometric characterization of the criticality of the preferred extremals. Sub-algebra of conformal symmetries consisting of generators for which conformal weight is integer multiple of given integer n is conjectured to act as critical deformations, that there are n conformal equivalence classes of extremals and that n defines the effective value of Planck constant $h_{eff} = n \times h$.

4.2 About Identification of the Preferred extremals of Kähler Action

Preferred extremal of Kähler action have remained one of the basic poorly defined notions of TGD. There are pressing motivations for understanding what the attribute “preferred” really means. Symmetries give a clue to the problem. The conformal invariance of string models naturally generalizes to 4-D invariance defined by quantum Yangian of quantum affine algebra (Kac-Moody type algebra) characterized by two complex coordinates and therefore explaining naturally the effective 2-dimensionality [K21]. Preferred extremal property should rely on this symmetry.

In Zero Energy Ontology (ZEO) preferred extremals are space-time surfaces connecting two space-like 3-surfaces at the ends of space-time surfaces at boundaries of causal diamond (CD). A natural looking condition is that the symplectic Noether charges associated with a sub-algebra of symplectic algebra with conformal weights n -multiples of the weights of the entire algebra vanish for preferred extremals. These conditions would be classical counterparts the the condition that super-symplectic sub-algebra annihilates the physical states. This would give a hierarchy of super-symplectic symmetry breakings and quantum criticalities having interpretation in terms of hierarchy of Planck constants $h_{eff} = n \times h$ identified as a hierarchy of dark matter. n could be interpreted as the number of space-time conformal gauge equivalence classes for space-time sheets connecting the 3-surfaces at the ends of space-time surface.

There are also many other proposals for what preferred extremal property could mean or imply. The weak form of electric-magnetic duality combined with the assumption that the contraction of the Kähler current with Kähler gauge potential vanishes for preferred extremals implies that Kähler action in Minkowskian space-time regions reduces to Chern-Simons terms at the light-like orbits of wormhole throats at which the signature of the induced metric changes its signature from Minkowskian to Euclidian. In regions with 4-D CP_2 projection (wormhole contacts) also a 3-D contribution not assignable to the boundary of the region might be possible. These conditions pose strong physically feasible conditions on extremals and might be true for preferred extremals too.

Number theoretic vision leads to a proposal that either the tangent space or normal space of given point of space-time surface is associative and thus quaternionic. Also the formulation in terms of quaternion holomorphy and quaternion-Kähler property is an attractive possibility. So called $M^8 - H$ duality is a variant of this vision and would mean that one can map associative/co-associative space-time surfaces from M^8 to H and also iterate this mapping from H to H to generate entire category of preferred extremals. The signature of M^4 is a general technical problem. For instance, the holomorphy in 2 complex variables could correspond to what I have called Hamilton-Jacobi property. Associativity/co-associativity of the tangent space makes sense also in Minkowskian signature.

In this chapter various views about preferred extremal property are discussed.

4.3 Construction of WCW Kähler Geometry from Symmetry Principles

There are three separate approaches to the challenge of constructing WCW Kähler geometry and spinor structure. The first one relies on a direct guess of the Kähler function. Second approach relies on the construction of Kähler form and metric utilizing the huge symmetries of the geometry needed to guarantee the mathematical existence of Riemann connection. The third approach relies on the construction of spinor structure assuming that complexified WCW gamma matrices are representable as linear combinations of fermionic oscillator operator for the second quantized free spinor fields at space-time surface and on the geometrization of super-conformal symmetries in terms of spinor structure.

In this chapter the construction of Kähler form and metric based on symmetries is discussed. The basic vision is that WCW can be regarded as the space of generalized Feynman diagrams with lines thickened to light-like 3-surfaces and vertices identified as partonic 2-surfaces. In zero energy ontology the strong form of General Coordinate Invariance (GCI) strongly suggests effective 2-dimensionality and the basic objects are taken to be pairs partonic 2-surfaces X^2 at opposite light-like boundaries of causal diamonds (CDs). This has however turned out to be too strong formulation for effective 2-dimensionality string world sheets carrying induced spinor fields are also present.

The hypothesis is that WCW can be regarded as a union of infinite-dimensional symmetric spaces G/H labeled by zero modes having an interpretation as classical, non-quantum fluctuating variables. A crucial role is played by the metric 2-dimensionality of the light-cone boundary δM_+^4 and of light-like 3-surfaces implying a generalization of conformal invariance. The group G acting as isometries of WCW is tentatively identified as the symplectic group of $\delta M_+^4 \times CP_2$. H corresponds to sub-group acting as diffeomorphisms at preferred 3-surface, which can be taken to correspond to maximum of Kähler function.

In zero energy ontology (ZEO) 3-surface corresponds to a pair of space-like 3-surfaces at the opposite boundaries of causal diamond (CD) and thus to a more or less unique extremal of Kähler action. The interpretation would be in terms of holography. One can also consider the inclusion of the light-like 3-surfaces at which the signature of the induced metric changes to the 3-surface so that it would become connected.

An explicit construction for the Hamiltonians of WCW isometry algebra as so called flux Hamiltonians using Hamiltonians of light-cone boundary is proposed and also the elements of Kähler form can be constructed in terms of these. Explicit expressions for WCW flux Hamiltonians as functionals of complex coordinates of the Cartesian product of the infinite-dimensional symmetric spaces having as points the partonic 2-surfaces defining the ends of the the light 3-surface (line of generalized Feynman diagram) are proposed.

This construction suffers from some rather obvious defects. Effective 2-dimensionality is realized in too strong sense, only covariantly constant right-handed neutrino is involved, and WCW Hamiltonians do not directly reflect the dynamics of Kähler action. The construction however generalizes in very straightforward manner to a construction free of these problems. This however requires the understanding of the dynamics of preferred extremals and Kähler-Dirac action.

4.4 WCW Spinor Structure

Quantum TGD should be reducible to the classical spinor geometry of the configuration space (“world of classical worlds” (WCW)). The possibility to express the components of WCW Kähler metric as anti-commutators of WCW gamma matrices becomes a practical tool if one assumes that WCW gamma matrices correspond to Noether super charges for super-symplectic algebra of WCW. The possibility to express the Kähler metric also in terms of Kähler function identified as Kähler for Euclidian space-time regions leads to a duality analogous to AdS/CFT duality.

Physical states should correspond to the modes of the WCW spinor fields and the identification of the fermionic oscillator operators as super-symplectic charges is highly attractive. WCW spinor fields cannot, as one might naively expect, be carriers of a definite spin and unit fermion number. Concerning the construction of the WCW spinor structure there are some important clues.

1. Geometrization of fermionic statistics in terms of WCW spinor structure

The great vision has been that the second quantization of the induced spinor fields can be under-

stood geometrically in terms of the WCW spinor structure in the sense that the anti-commutation relations for WCW gamma matrices require anti-commutation relations for the oscillator operators for free second quantized induced spinor fields.

1. One must identify the counterparts of second quantized fermion fields as objects closely related to the WCW spinor structure. Ramond model has as its basic field the anti-commuting field $\Gamma^k(x)$, whose Fourier components are analogous to the gamma matrices of the WCW and which behaves like a spin 3/2 fermionic field rather than a vector field. This suggests that the complexified gamma matrices of the WCW are analogous to spin 3/2 fields and therefore expressible in terms of the fermionic oscillator operators so that their anti-commutativity naturally derives from the anti-commutativity of the fermionic oscillator operators.

As a consequence, WCW spinor fields can have arbitrary fermion number and there would be hopes of describing the whole physics in terms of WCW spinor field. Clearly, fermionic oscillator operators would act in degrees of freedom analogous to the spin degrees of freedom of the ordinary spinor and bosonic oscillator operators would act in degrees of freedom analogous to the “orbital” degrees of freedom of the ordinary spinor field.

2. The classical theory for the bosonic fields is an essential part of the WCW geometry. It would be very nice if the classical theory for the spinor fields would be contained in the definition of the WCW spinor structure somehow. The properties of the modified massless Dirac operator associated with the induced spinor structure are indeed very physical. The modified massless Dirac equation for the induced spinors predicts a separate conservation of baryon and lepton numbers. The differences between quarks and leptons result from the different couplings to the CP_2 Kähler potential. In fact, these properties are shared by the solutions of massless Dirac equation of the imbedding space.
3. Since TGD should have a close relationship to the ordinary quantum field theories it would be highly desirable that the second quantized free induced spinor field would somehow appear in the definition of the WCW geometry. This is indeed true if the complexified WCW gamma matrices are linearly related to the oscillator operators associated with the second quantized induced spinor field on the space-time surface and/or its boundaries. There is actually no deep reason forbidding the gamma matrices of the WCW to be spin half odd-integer objects whereas in the finite-dimensional case this is not possible in general. In fact, in the finite-dimensional case the equivalence of the spinorial and vectorial vielbeins forces the spinor and vector representations of the vielbein group $SO(D)$ to have same dimension and this is possible for $D = 8$ -dimensional Euclidian space only. This coincidence might explain the success of 10-dimensional super string models for which the physical degrees of freedom effectively correspond to an 8-dimensional Euclidian space.
4. It took a long time to realize that the ordinary definition of the gamma matrix algebra in terms of the anti-commutators $\{\gamma_A, \gamma_B\} = 2g_{AB}$ must in TGD context be replaced with $\{\gamma_A^\dagger, \gamma_B\} = iJ_{AB}$, where J_{AB} denotes the matrix elements of the Kähler form of the WCW. The presence of the Hermitian conjugation is necessary because WCW gamma matrices carry fermion number. This definition is numerically equivalent with the standard one in the complex coordinates. The realization of this delicacy is necessary in order to understand how the square of the WCW Dirac operator comes out correctly.

2. Kähler-Dirac equation for induced spinor fields

Super-symmetry between fermionic and and WCW degrees of freedom dictates that Kähler-Dirac action is the unique choice for the Dirac action

There are several approaches for solving the Kähler-Dirac (or Kähler-Dirac) equation.

1. The most promising approach assumes that the solutions are restricted on 2-D stringy world sheets and/or partonic 2-surfaces. This strange looking view is a rather natural consequence of both strong form of holography and of number theoretic vision, and also follows from the notion of finite measurement resolution having discretization at partonic 2-surfaces as a geometric correlate. Furthermore, the conditions stating that electric charge is well-defined

for preferred extremals forces the localization of the modes to 2-D surfaces in the generic case. This also resolves the interpretational problems related to possibility of strong parity breaking effects since induce W fields and possibly also Z^0 field above weak scale, vanish at these surfaces.

The condition that also spinor dynamics is associative suggests strongly that the localization to 2-D surface occurs always (for right-handed neutrino the above conditions does not apply this). The induced gauge potentials are the possible source of trouble but the holomorphy of spinor modes completely analogous to that encountered in string models saves the situation. Whether holomorphy could be replaced with its quaternionic counterpart in Euclidian regions is not clear (this if W fields vanish at the entire space-time surface so that 4-D modes are possible). Neither it is clear whether the localization to 2-D surfaces occurs also in Euclidian regions with 4-D CP_2 projection.

2. One expects that stringy approach based on 4-D generalization of conformal invariance or its 2-D variant at 2-D preferred surfaces should also allow to understand the Kähler-Dirac equation. Conformal invariance indeed allows to write the solutions explicitly using formulas similar to encountered in string models. In accordance with the earlier conjecture, all modes of the Kähler-Dirac operator generate badly broken super-symmetries.
3. Well-definedness of em charge is not enough to localize spinor modes at string world sheets. Covariantly constant right-handed neutrino certainly defines solutions de-localized inside entire space-time sheet. This need not be the case if right-handed neutrino is not covariantly constant since the non-vanishing CP_2 part for the induced gamma matrices mixes it with left-handed neutrino. For massless extremals (at least) the CP_2 part however vanishes and right-handed neutrino allows also massless holomorphic modes de-localized at entire space-time surface and the de-localization inside Euclidian region defining the line of generalized Feynman diagram is a good candidate for the right-handed neutrino generating the least broken super-symmetry. This super-symmetry seems however to differ from the ordinary one in that ν_R is expected to behave like a passive spectator in the scattering. Also for the left-handed neutrino solutions localized inside string world sheet the condition that coupling to right-handed neutrino vanishes is guaranteed if gamma matrices are either purely Minkowskian or CP_2 like inside the world sheet.

awcwspin

Quantum TGD should be reducible to the classical spinor geometry of the configuration space (“world of classical worlds” (WCW)). The possibility to express the components of WCW Kähler metric as anti-commutators of WCW gamma matrices becomes a practical tool if one assumes that WCW gamma matrices correspond to Noether super charges for super-symplectic algebra of WCW. The possibility to express the Kähler metric also in terms of Kähler function identified as Kähler for Euclidian space-time regions leads to a duality analogous to AdS/CFT duality.

Physical states should correspond to the modes of the WCW spinor fields and the identification of the fermionic oscillator operators as super-symplectic charges is highly attractive. WCW spinor fields cannot, as one might naively expect, be carriers of a definite spin and unit fermion number. Concerning the construction of the WCW spinor structure there are some important clues.

1. Geometrization of fermionic statistics in terms of WCW spinor structure

The great vision has been that the second quantization of the induced spinor fields can be understood geometrically in terms of the WCW spinor structure in the sense that the anti-commutation relations for WCW gamma matrices require anti-commutation relations for the oscillator operators for free second quantized induced spinor fields.

1. One must identify the counterparts of second quantized fermion fields as objects closely related to the WCW spinor structure. Ramond model has as its basic field the anti-commuting field $\Gamma^k(x)$, whose Fourier components are analogous to the gamma matrices of the WCW and which behaves like a spin 3/2 fermionic field rather than a vector field. This suggests that the complexified gamma matrices of the WCW are analogous to spin 3/2 fields and therefore expressible in terms of the fermionic oscillator operators so that their anti-commutativity naturally derives from the anti-commutativity of the fermionic oscillator operators.

As a consequence, WCW spinor fields can have arbitrary fermion number and there would be hopes of describing the whole physics in terms of WCW spinor field. Clearly, fermionic oscillator operators would act in degrees of freedom analogous to the spin degrees of freedom of the ordinary spinor and bosonic oscillator operators would act in degrees of freedom analogous to the “orbital” degrees of freedom of the ordinary spinor field.

2. The classical theory for the bosonic fields is an essential part of the WCW geometry. It would be very nice if the classical theory for the spinor fields would be contained in the definition of the WCW spinor structure somehow. The properties of the modified massless Dirac operator associated with the induced spinor structure are indeed very physical. The modified massless Dirac equation for the induced spinors predicts a separate conservation of baryon and lepton numbers. The differences between quarks and leptons result from the different couplings to the CP_2 Kähler potential. In fact, these properties are shared by the solutions of massless Dirac equation of the imbedding space.
3. Since TGD should have a close relationship to the ordinary quantum field theories it would be highly desirable that the second quantized free induced spinor field would somehow appear in the definition of the WCW geometry. This is indeed true if the complexified WCW gamma matrices are linearly related to the oscillator operators associated with the second quantized induced spinor field on the space-time surface and/or its boundaries. There is actually no deep reason forbidding the gamma matrices of the WCW to be spin half odd-integer objects whereas in the finite-dimensional case this is not possible in general. In fact, in the finite-dimensional case the equivalence of the spinorial and vectorial vielbeins forces the spinor and vector representations of the vielbein group $SO(D)$ to have same dimension and this is possible for $D = 8$ -dimensional Euclidian space only. This coincidence might explain the success of 10-dimensional super string models for which the physical degrees of freedom effectively correspond to an 8-dimensional Euclidian space.
4. It took a long time to realize that the ordinary definition of the gamma matrix algebra in terms of the anti-commutators $\{\gamma_A, \gamma_B\} = 2g_{AB}$ must in TGD context be replaced with $\{\gamma_A^\dagger, \gamma_B\} = iJ_{AB}$, where J_{AB} denotes the matrix elements of the Kähler form of the WCW. The presence of the Hermitian conjugation is necessary because WCW gamma matrices carry fermion number. This definition is numerically equivalent with the standard one in the complex coordinates. The realization of this delicacy is necessary in order to understand how the square of the WCW Dirac operator comes out correctly.

2. Kähler-Dirac equation for induced spinor fields

Super-symmetry between fermionic and and WCW degrees of freedom dictates that Kähler-Dirac action is the unique choice for the Dirac action

There are several approaches for solving the Kähler-Dirac (or Kähler-Dirac) equation.

1. The most promising approach assumes that the solutions are restricted on 2-D stringy world sheets and/or partonic 2-surfaces. This strange looking view is a rather natural consequence of both strong form of holography and of number theoretic vision, and also follows from the notion of finite measurement resolution having discretization at partonic 2-surfaces as a geometric correlate. Furthermore, the conditions stating that electric charge is well-defined for preferred extremals forces the localization of the modes to 2-D surfaces in the generic case. This also resolves the interpretational problems related to possibility of strong parity breaking effects since induce W fields and possibly also Z^0 field above weak scale, vanish at these surfaces.

The condition that also spinor dynamics is associative suggests strongly that the localization to 2-D surface occurs always (for right-handed neutrino the above conditions does not apply this). The induced gauge potentials are the possible source of trouble but the holomorphy of spinor modes completely analogous to that encountered in string models saves the situation. Whether holomorphy could be replaced with its quaternionic counterpart in Euclidian regions is not clear (this if W fields vanish at the entire space-time surface so that 4-D modes are possible). Neither it is clear whether the localization to 2-D surfaces occurs also in Euclidian regions with 4-D CP_2 projection.

2. One expects that stringy approach based on 4-D generalization of conformal invariance or its 2-D variant at 2-D preferred surfaces should also allow to understand the Kähler-Dirac equation. Conformal invariance indeed allows to write the solutions explicitly using formulas similar to encountered in string models. In accordance with the earlier conjecture, all modes of the Kähler-Dirac operator generate badly broken super-symmetries.
3. Well-definedness of em charge is not enough to localize spinor modes at string world sheets. Covariantly constant right-handed neutrino certainly defines solutions de-localized inside entire space-time sheet. This need not be the case if right-handed neutrino is not covariantly constant since the non-vanishing CP_2 part for the induced gamma matrices mixes it with left-handed neutrino. For massless extremals (at least) the CP_2 part however vanishes and right-handed neutrino allows also massless holomorphic modes de-localized at entire space-time surface and the de-localization inside Euclidian region defining the line of generalized Feynman diagram is a good candidate for the right-handed neutrino generating the least broken super-symmetry. This super-symmetry seems however to differ from the ordinary one in that ν_R is expected to behave like a passive spectator in the scattering. Also for the left-handed neutrino solutions localized inside string world sheet the condition that coupling to right-handed neutrino vanishes is guaranteed if gamma matrices are either purely Minkowskian or CP_2 like inside the world sheet.

4.5 Recent View about Kähler Geometry and Spin Structure of WCW

The construction of Kähler geometry of WCW (“world of classical worlds”) is fundamental to TGD program. I ended up with the idea about physics as WCW geometry around 1985 and made a breakthrough around 1990, when I realized that Kähler function for WCW could correspond to Kähler action for its preferred extremals defining the analogs of Bohr orbits so that classical theory with Bohr rules would become an exact part of quantum theory and path integral would be replaced with genuine integral over WCW. The motivating construction was that for loop spaces leading to a unique Kähler geometry. The geometry for the space of 3-D objects is even more complex than that for loops and the vision still is that the geometry of WCW is unique from the mere existence of Riemann connection.

This chapter represents the updated version of the construction providing a solution to the problems of the previous construction. The basic formulas remain as such but the expressions for WCW super-Hamiltonians defining WCW Hamiltonians (and matrix elements of WCW metric) as their anticommutator are replaced with those following from the dynamics of the Kähler-Dirac action.

4.6 The Classical Part of the Twistor Story

Twistor Grassmannian formalism has made a breakthrough in $\mathcal{N} = 4$ supersymmetric gauge theories and the Yangian symmetry suggests that much more than mere technical breakthrough is in question. Twistors seem to be tailor made for TGD but it seems that the generalization of twistor structure to that for 8-D imbedding space $H = M^4 \times CP_2$ is necessary. M^4 (and S^4 as its Euclidian counterpart) and CP_2 are indeed unique in the sense that they are the only 4-D spaces allowing twistor space with Kähler structure.

The Cartesian product of twistor spaces $P_3 = SU(2, 2)/SU(2, 1) \times U(1)$ and F_3 defines twistor space for the imbedding space H and one can ask whether this generalized twistor structure could allow to understand both quantum TGD and classical TGD defined by the extremals of Kähler action. In the following I summarize the background and develop a proposal for how to construct extremals of Kähler action in terms of the generalized twistor structure. One ends up with a scenario in which space-time surfaces are lifted to twistor spaces by adding CP_1 fiber so that the twistor spaces give an alternative representation for generalized Feynman diagrams.

There is also a very closely analogy with superstring models. Twistor spaces replace Calabi-Yau manifolds and the modification recipe for Calabi-Yau manifolds by removal of singularities can be applied to remove self-intersections of twistor spaces and mirror symmetry emerges naturally. The overall important implication is that the methods of algebraic geometry used in super-string theories should apply in TGD framework.

The physical interpretation is totally different in TGD. The landscape is replaced with twistor spaces of space-time surfaces having interpretation as generalized Feynman diagrams and twistor spaces as sub-manifolds of $P_3 \times F_3$ replace Witten's twistor strings.

The classical view about twistorialization of TGD makes possible a more detailed formulation of the previous ideas about the relationship between TGD and Witten's theory and twistor Grassmann approach. Furthermore, one ends up to a formulation of the scattering amplitudes in terms of Yangian of the super-symplectic algebra relying on the idea that scattering amplitudes are sequences consisting of algebraic operations (product and co-product) having interpretation as vertices in the Yangian extension of super-symplectic algebra. These sequences connect given initial and final states and having minimal length. One can say that Universe performs calculations.

4.7 Unified Number Theoretical Vision

An updated view about $M^8 - H$ duality is discussed. $M^8 - H$ duality allows to deduce $M^4 \times CP_2$ via number theoretical compactification. One important correction is that octonionic spinor structure makes sense only for M^8 whereas for $M^4 \times CP_2$ complexified quaternions characterized the spinor structure.

Octonions, quaternions associative and co-associative space-time surfaces, octonionic spinors and twistors and twistor spaces are highly relevant for quantum TGD. In the following some general observations distilled during years are summarized.

There is a beautiful pattern present suggesting that $H = M^4 \times CP_2$ is completely unique on number theoretical grounds. Consider only the following facts. M^4 and CP_2 are the unique 4-D spaces allowing twistor space with Kähler structure. Octonionic projective space OP_2 appears as octonionic twistor space (there are no higher-dimensional octonionic projective spaces). Octotwistors generalise the twistorial construction from M^4 to M^8 and octonionic gamma matrices make sense also for H with quaternionicity condition reducing OP_2 to 12-D $G_2/U(1) \times U(1)$ having same dimension as the twistor space $CP_3 \times SU(3)/U(1) \times U(1)$ of H assignable to complexified quaternionic representation of gamma matrices.

A further fascinating structure related to octo-twistors is the non-associated analog of Lie group defined by automorphisms by octonionic imaginary units: this group is topologically six-sphere. Also the analogy of quaternionicity of preferred extremals in TGD with the Majorana condition central in super string models is very thought provoking. All this suggests that associativity indeed could define basic dynamical principle of TGD.

Number theoretical vision about quantum TGD involves both p-adic number fields and classical number fields and the challenge is to unify these approaches. The challenge is non-trivial since the p-adic variants of quaternions and octonions are not number fields without additional conditions. The key idea is that TGD reduces to the representations of Galois group of algebraic numbers realized in the spaces of octonionic and quaternionic adeles generalizing the ordinary adeles as Cartesian products of all number fields: this picture relates closely to Langlands program. Associativity would force sub-algebras of the octonionic adeles defining 4-D surfaces in the space of octonionic adeles so that 4-D space-time would emerge naturally. $M^8 - H$ correspondence in turn would map the space-time surface in M^8 to $M^4 \times CP_2$.

A long-standing question has been the origin of preferred p-adic primes characterizing elementary particles. I have proposed several explanations and the most convincing hitherto is related to the algebraic extensions of rationals and p-adic numbers selecting naturally preferred primes as those which are ramified for the extension in question.

4.8 Knots and TGD

Khovanov homology generalizes the Jones polynomial as knot invariant. The challenge is to find a quantum physical construction of Khovanov homology analogous to the topological QFT defined by Chern-Simons action allowing to interpret Jones polynomial as vacuum expectation value of Wilson loop in non-Abelian gauge theory.

Witten's approach to Khovanov homology relies on fivebranes as is natural if one tries to define 2-knot invariants in terms of their cobordisms involving violent un-knottings. Despite the difference in approaches it is very useful to try to find the counterparts of this approach in quantum TGD since this would allow to gain new insights to quantum TGD itself as almost topological QFT

identified as symplectic theory for 2-knots, braids and braid cobordisms. This comparison turns out to be extremely useful from TGD point of view.

1. Key question concerns the identification of string world sheets. A possible identification of string world sheets and therefore also of the braids whose ends carry quantum numbers of many particle states at partonic 2-surfaces emerges if one identifies the string world sheets as singular surfaces in the same manner as is done in Witten's approach.

In TGD framework the localization of the modes of the induced spinor fields at 2-D surfaces carrying vanishing induced W boson fields guaranteeing that the em charge of spinor modes is well-defined for a generic preferred extremal is natural. Besides string world sheets partonic 2-surfaces are good candidates for this kind of surfaces. It is not clear whether one can have continuous slicing of this kind by string world sheets and partonic 2-surfaces orthogonal to them or whether only discrete set of these surfaces is possible.

2. Also a physical interpretation of the operators Q , F , and P of Khovanov homology emerges. P would correspond to instanton number and F to the fermion number assignable to right handed neutrinos. The breaking of M^4 chiral invariance makes possible to realize Q physically. The finding that the generalizations of Wilson loops can be identified in terms of the gerbe fluxes $\int H_A J$ supports the conjecture that TGD as almost topological QFT corresponds essentially to a symplectic theory for braids and 2-knots.

The basic challenge of quantum TGD is to give a precise content to the notion of generalized Feynman diagram and the reduction to braids of some kind is very attractive possibility inspired by zero energy ontology. The point is that no $n > 2$ -vertices at the level of braid strands are needed if bosonic emergence holds true.

1. For this purpose the notion of algebraic knot is introduced and the possibility that it could be applied to generalized Feynman diagrams is discussed. The algebraic structures kei , quandle, rack, and biquandle and their algebraic modifications as such are not enough. The lines of Feynman graphs are replaced by braids and in vertices braid strands redistribute. This poses several challenges: the crossing associated with braiding and crossing occurring in non-planar Feynman diagrams should be integrated to a more general notion; braids are replaced with sub-manifold braids; braids of braids ...of braids are possible; the redistribution of braid strands in vertices should be algebraized. In the following I try to abstract the basic operations which should be algebraized in the case of generalized Feynman diagrams.
2. One should be also able to concretely identify braids and 2-braids (string world sheets) as well as partonic 2-surfaces and I have discussed several identifications during last years. Legendrian braids turn out to be very natural candidates for braids and their duals for the partonic 2-surfaces. String world sheets in turn could correspond to the analogs of Lagrangian sub-manifolds or two minimal surfaces of space-time surface satisfying the weak form of electric-magnetic duality. The latter option turns out to be more plausible. This identification - if correct - would solve quantum TGD explicitly at string world sheet level which corresponds to finite measurement resolution.
3. Also a brief summary of generalized Feynman rules in zero energy ontology is proposed. This requires the identification of vertices, propagators, and prescription for integrating over all 3-surfaces. It turns out that the basic building blocks of generalized Feynman diagrams are well-defined.
4. The notion of generalized Feynman diagram leads to a beautiful duality between the descriptions of hadronic reactions in terms of hadrons and partons analogous to gauge-gravity duality and AdS/CFT duality but requiring no additional assumptions. The model of quark gluon plasma as a strongly interacting phase is proposed. Color magnetic flux tubes are responsible for the long range correlations making the plasma phase more like a very large hadron rather than a gas of partons. One also ends up with a simple estimate for the viscosity/entropy ratio using black-hole analogy.

REFERENCES

Mathematics

- [A1] Pope CN. Eigenfunctions and $Spin^c$ Structures on CP_2 , 1980.
- [A2] Hanson J Eguchi T, Gilkey B. *Phys Rep*, 66:1980, 1980.
- [A3] Eisenhart. *Riemannian Geometry*. Princeton University Press, 1964.
- [A4] Pope CN Gibbons GW. CP_2 as gravitational instanton. *Comm Math Phys*, 55, 1977.
- [A5] Pope CN Hawking SW. Generalized Spin Structures in Quantum Gravity. *Phys Lett*, (1), 1978.
- [A6] Milnor J. *Topology form Differential Point of View*. The University Press of Virginia, Virginia, 1965.
- [A7] Spivak M. *Differential Geometry I,II,III,IV*. Publish or Perish, Boston, 1970.
- [A8] Thom R. *Comm Math Helvet*, 28, 1954.
- [A9] Wallace. *Differential Topology*. W. A. Benjamin, New York, 1968.

Cosmology and Astro-Physics

- [E1] Nottale L Da Rocha D. Gravitational Structure Formation in Scale Relativity. Available at: <http://arxiv.org/abs/astro-ph/0310036>, 2003.

Books related to TGD

- [K1] Pitkänen M. *Topological Geometro-dynamics*. 1983.
- [K2] Pitkänen M. *Bio-Systems as Conscious Holograms*. Onlinebook. Available at: http://tgdtheory.fi/public_html/hologram/hologram.html, 2006.
- [K3] Pitkänen M. *Bio-Systems as Self-Organizing Quantum Systems*. Onlinebook. Available at: http://tgdtheory.fi/public_html/bioselforg/bioselforg.html, 2006.
- [K4] Pitkänen M. Does TGD Predict the Spectrum of Planck Constants? In *Hyper-finite Factors and Dark Matter Hierarchy*. Onlinebook. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html#Planck, 2006.
- [K5] Pitkänen M. *Genes and Memes*. Onlinebook. Available at: http://tgdtheory.fi/public_html/genememe/genememe.html, 2006.
- [K6] Pitkänen M. *Magnetospheric Consciousness*. Onlinebook. Available at: http://tgdtheory.fi/public_html/magnconsc/magnconsc.html, 2006.
- [K7] Pitkänen M. *Mathematical Aspects of Consciousness Theory*. Onlinebook. Available at: http://tgdtheory.fi/public_html/mathconsc/mathconsc.html, 2006.
- [K8] Pitkänen M. *Physics in Many-Sheeted Space-Time*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html, 2006.
- [K9] Pitkänen M. *Quantum Hardware of Living Matter*. Onlinebook. Available at: http://tgdtheory.fi/public_html/bioware/bioware.html, 2006.
- [K10] Pitkänen M. Quantum Model for Bio-Superconductivity: I. In *TGD and EEG*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdeeg/tgdeeg.html#biosupercondI, 2006.

- [K11] Pitkänen M. Quantum Model for Bio-Superconductivity: II. In *TGD and EEG*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdeeg/tgdeeg.html#biosupercondII, 2006.
- [K12] Pitkänen M. *Quantum Physics as Infinite-Dimensional Geometry*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdgeom/tgdgeom.html, 2006.
- [K13] Pitkänen M. TGD and Astrophysics. In *Physics in Many-Sheeted Space-Time*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdclass/tgdclass.html#astro, 2006.
- [K14] Pitkänen M. *TGD and EEG*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdeeg/tgdeeg.html, 2006.
- [K15] Pitkänen M. *TGD and Fringe Physics*. Onlinebook. Available at: http://tgdtheory.fi/public_html/freenergy/freenergy.html, 2006.
- [K16] Pitkänen M. *TGD as a Generalized Number Theory*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html, 2006.
- [K17] Pitkänen M. TGD as a Generalized Number Theory: Quaternions, Octonions, and their Hyper Counterparts. In *TGD as a Generalized Number Theory*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#visionb, 2006.
- [K18] Pitkänen M. *TGD Inspired Theory of Consciousness*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdconsc/tgdconsc.html, 2006.
- [K19] Pitkänen M. The classical part of the twistor story. In *Towards M-Matrix*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#twistorstory, 2006.
- [K20] Pitkänen M. *Topological Geometrodynamics: an Overview*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdview/tgdview.html, 2006.
- [K21] Pitkänen M. Yangian Symmetry, Twistors, and TGD. In *Towards M-Matrix*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#Yangian, 2006.
- [K22] Pitkänen M. Construction of Quantum Theory: More about Matrices. In *Towards M-Matrix*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html#UandM, 2012.
- [K23] Pitkänen M. Quantum Mind and Neuroscience. In *TGD based view about living matter and remote mental interactions*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdlian/tgdlian.html#lianPN, 2012.
- [K24] Pitkänen M. *TGD Based View About Living Matter and Remote Mental Interactions*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdlian/tgdlian.html, 2012.
- [K25] Pitkänen M. *Hyper-finite Factors and Dark Matter Hierarchy*. Onlinebook. Available at: http://tgdtheory.fi/public_html/neuplanck/neuplanck.html, 2013.
- [K26] Pitkänen M. *p-Adic length Scale Hypothesis*. Onlinebook. Available at: http://tgdtheory.fi/public_html/padphys/padphys.html, 2013.
- [K27] Pitkänen M. *Quantum TGD*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdquantum/tgdquantum.html, 2013.
- [K28] Pitkänen M. What p-Adic Icosahedron Could Mean? And What about p-Adic Manifold? In *TGD as a Generalized Number Theory*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#picosahedron, 2013.
- [K29] Pitkänen M. Unified Number Theoretical Vision. In *TGD as a Generalized Number Theory*. Onlinebook. Available at: http://tgdtheory.fi/public_html/tgdnumber/tgdnumber.html#numbervision, 2014.

Articles about TGD

[L1] Pitkänen M. CMAP representations about TGD. Available at: <http://www.tgdtheory.fi/cmaphtml.html>, 2014.

[L2] Pitkänen M. CMAP representations about TGD, and TGD inspired theory of consciousness and quantum biology. Available at: <http://www.tgdtheory.fi/tgdglossary.pdf>, 2014.